

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
14 February 2002 (14.02.2002)

PCT

(10) International Publication Number  
WO 02/11669 A2

- (51) International Patent Classification<sup>7</sup>: A61K (74) Agents: ANTILER, Adriane, M., et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).
- (21) International Application Number: PCT/US01/23098
- (22) International Filing Date: 20 July 2001 (20.07.2001) (81) Designated States (national): AU, CA, JP.
- (25) Filing Language: English (84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).
- (26) Publication Language: English
- (30) Priority Data: 60/223,133 7 August 2000 (07.08.2000) US Published:  
— without international search report and to be republished upon receipt of that report
- (71) Applicant: ANTIGENICS, LLC [US/US]; Suite 2100, 630 Fifth Avenue, New York, NY 10011 (US).
- (72) Inventor: ARMEN, Garo, H.; 66 Mayfair Lane, Manhasset, NY 11030 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/11669 A2

(54) Title: COMPOSITIONS COMPRISING HEAT SHOCK PROTEINS OR ALPHA(2)MACROGLOBULIN, ANTIGENIC MOLECULES AND SAPONINS, AND METHODS OF USE THEREOF

(57) Abstract: The present invention relates to pharmaceutical compositions and methods for the prevention and treatment of autoimmune diseases, infectious diseases, neurodegenerative diseases, and primary and metastatic neoplastic diseases. In the practice of the invention, the compositions are employed comprising: (a) a heat shock protein (hsp) or an alpha(2)macroglobulin ( $\alpha$ 2M); (b) a saponin; and, optionally, (c) an antigenic molecule. The antigenic molecule displays the antigenicity of an antigen of: (a) a cell that elicits an autoimmune response; (b) an agent of an infectious disease; (c) a cancerous cell; or (d) a cell or structure associated with a neurodegenerative or amyloid disease. The hsp's that can be used in the practice of the invention include but are not limited to hsp70, hsp90, gp96, calreticulin, hsp 110, grp170, and PDI, alone or in combination with each other. The antigenic molecule can be covalently or noncovalently bound to the hsp or  $\alpha$ 2M, free in solution, and/or covalently bound to the saponin. The compositions of the invention can be administered alone or in combination with the administration or antigen presenting cells sensitized with an hsp- or  $\alpha$ 2M-antigenic molecule complex.

COMPOSITIONS COMPRISING HEAT SHOCK PROTEINS OR ALPHA(2)MACROGLOBULIN,  
ANTIGENIC MOLECULES AND SAPONINS, AND METHODS OF USE THEREOF

---

**1. Field of the Invention**

5           The present invention relates to pharmaceutical compositions that are useful for the prevention and treatment of infectious diseases, primary and metastatic neoplastic diseases (*i.e.*, cancer), neurodegenerative or amyloid diseases, and autoimmune diseases, and methods of formulating the compositions. The compositions comprise a heat shock protein (hsp) or alpha(2)macroglobulin ( $\alpha$ 2M) and a saponin when used for the treatment  
10 and prevention of an autoimmune disease. The compositions further comprise an antigenic molecule when used for the treatment or prevention of cancer, infectious disease or neurodegenerative or amyloid disease. Compositions comprising antigenic molecules may also be used for the treatment or prevention of autoimmune disorders.

**2. Background of the Invention**

**2.1. Vaccines**

Vaccination has eradicated certain diseases such as polio, tetanus, chicken pox, and measles in many countries. This approach has exploited the ability of the immune system to resist and prevent infectious diseases.

20           Traditional ways of preparing vaccines include the use of inactivated or attenuated pathogens. A suitable inactivation of the pathogenic microorganism renders it harmless as a biological agent but does not destroy its immunogenicity. Injection of these "killed" particles into a host will then elicit an immune response capable of preventing a future infection with a live microorganism. However, a major concern in the use of  
25 inactivated pathogens as vaccines is the failure to inactivate all the microorganisms. Even when complete microorganism inactivation is accomplished, the immunity achieved is still often incomplete and/or short-lived, requiring multiple immunizations to sustain an immune response (presumably since killed pathogens do not multiply in their host or for other unknown reasons). Finally, the inactivation process may alter the microorganism's  
30 antigens, rendering them less effective as immunogens.

Attenuation refers to the production of strains of pathogenic microorganisms which have essentially lost their disease-producing ability. One way to accomplish this is to subject the microorganism to unusual growth conditions and/or frequent passage in cell culture. Mutants are then selected which have lost virulence but yet are capable of eliciting  
35 an immune response. Attenuated pathogens often make good immunogens as they actually replicate in the host cell and elicit long-lasting immunity. However, several problems are

encountered with the use of live vaccines, the most worrisome being insufficient attenuation and the risk of reversion to virulence.

An alternative to the above methods is the use of subunit vaccines. Subunit vaccines involve immunization only with a microorganism's components which contain the relevant immunological material. Another promising alternative is the use of DNA or RNA as vaccines. Such genetic vaccines have progressed from idea to reality and are being investigated in clinical trials (*see*, Weiner and Kennedy, July 1999, Scientific American, pp. 50-57).

Vaccines are often formulated to contain various adjuvants in addition to antigen. Adjuvants aid in attaining a more durable and higher level of immunity using smaller amounts of antigen or fewer doses than if the antigen were administered alone. However, the mechanism(s) of adjuvants is not completely understood and is still unpredictable (*see* Suzue *et al.*, 1996, Basel: Birkhauser Verlag, 454-55).

## 2.2. Immune Responses

An organism's immune system responds to pathogens or other harmful agents by two basic mechanisms – a humoral response and a cell-mediated response (*see* Alberts *et al.*, 1994, Molecular Biology of the Cell, pp. 1195-96). When resting B cells are activated by antigen to proliferate and mature into antibody-secreting cells, they produce and secrete antibodies with a unique antigen-binding site. This antibody-secreting reaction is known as the humoral response. On the other hand, the diverse responses of T cells are collectively called cell-mediated immune reactions. There are two main classes of T cells – cytotoxic T cells and helper T cells. Cytotoxic T cells directly kill cells that are infected with a virus or some other intracellular microorganism. Helper T cells, by contrast, help stimulate the responses of other cells: they help activate macrophages, dendritic cells and B cells, for example (Alberts *et al.*, 1994, Molecular Biology of the Cell, p. 1228). Both cytotoxic T cells and helper T cells recognize antigen in the form of peptide fragments that are generated by the degradation of foreign protein antigens inside the target cell, and both, therefore, depend on major histocompatibility complex (MHC) molecules, which bind these peptide fragments, carry them to the cell surface, and present them there to the T cells (Alberts *et al.*, *Id.*). MHC molecules are typically found in abundance on antigen-presenting cells (APCs).

Srivastava *et al.* demonstrated immune response to methylcholanthrene-induced sarcomas of inbred mice (1988, Immunol. Today 9:78-83). In these studies, it was found that the molecules responsible for the individually distinct immunogenicity of these tumors were glycoproteins of 96kDa (gp96) and intracellular proteins of 84 to 86kDa

- (Srivastava *et al.*, 1986, Proc. Natl. Acad. Sci. USA 83:3407-3411; Ullrich *et al.*, 1986, Proc. Natl. Acad. Sci. USA 83:3121-3125). Immunization of mice with gp96 or p84/86 isolated from a particular tumor rendered the mice immune to that particular tumor, but not to antigenically distinct tumors. Isolation and characterization of genes encoding gp96 and p84/86 revealed significant homology between them, and showed that gp96 and p84/86 were, respectively, the endoplasmic reticular and cytosolic counterparts of the same heat shock proteins (Srivastava *et al.*, 1988, Immunogenetics 28:205-207; Srivastava *et al.*, 1991, Curr. Top. Microbiol. Immunol. 167:109-123). Further, hsp70 was shown to elicit immunity to the tumor from which it was isolated but not to antigenically distinct tumors. However, hsp70 depleted of peptides was found to lose its immunogenic activity (Udono and Srivastava, 1993, J. Exp. Med. 178:1391-1396). These observations suggested that the heat shock proteins are not immunogenic *per se*, but instead form noncovalent complexes with antigenic peptides, and the complexes elicit specific immunity to the antigenic peptides (Srivastava, 1993, Adv. Cancer Res. 62:153-177; Udono *et al.*, 1994, J. Immunol., 152:5398-5403; Suto *et al.*, 1995, Science, 269:1585-1588).

- Noncovalent complexes of hsps and peptide, purified from cancer cells, can be used for the treatment and prevention of cancer and have been described in PCT publications WO 96/10411, dated April 11, 1996, and WO 97/10001, dated March 20, 1997 (U.S. Patent No. 5,750,119 issued April 12, 1998, and U.S. Patent No. 5,837,251 issued November 17, 1998, respectively, each of which is incorporated by reference herein in its entirety). The isolation and purification of stress protein-peptide complexes has been described, for example, from pathogen-infected cells, and can be used for the treatment and prevention of infection caused by pathogens such as viruses and other intracellular pathogens, including bacteria, protozoa, fungi and parasites (*see e.g.*, PCT Publication WO 95/24923, dated September 21, 1995). Immunogenic stress protein-peptide complexes can also be prepared by *in vitro* complexing of stress protein and antigenic peptides, and the uses of such complexes for the treatment and prevention of cancer and infectious diseases has been described in PCT publication WO 97/10000, dated March 20, 1997 and U.S. Patent No. 6,030,618 issued February 29, 2000. The use of stress protein-peptide complexes for sensitizing antigen presenting cells *in vitro* for use in adoptive immunotherapy is described in PCT publication WO 97/10002, dated March 20, 1997 (*see also*, U.S. Patent No. 5,985,270 issued November 16, 1999).

### 2.3. Heat Shock Proteins and Their Roles in Antigen Presentation

#### 2.3.1. Heat Shock Proteins

Heat shock proteins (hsps), also referred to as stress proteins, were first identified as proteins synthesized by cells in response to heat shock. hsps have been  
5 classified into five families based on molecular weight, *i.e.* hsp100, hsp90, hsp70, hsp60, and smhsp. Many members of these families were found subsequently to be induced in response to other stressful stimuli, including nutrient deprivation, metabolic disruption, oxygen radicals, and infection with intracellular pathogens (*see* Welch, May 1993, Scientific American 56-64; Young, 1990, Annu. Rev. Immunol. 8:401-420; Craig, 1993,  
10 Science 260:1902-1903; Gething *et al.*, 1992, Nature 355:33-45; and Lindquist *et al.*, 1988, Annu. Rev. Genetics 22:631-677).

Heat shock proteins are among the most highly conserved proteins in existence. For example, DnaK, the hsp70 from *E. coli*, has about 50% amino acid sequence  
15 identity with hsp70 proteins from excoriates (Bardwell *et al.*, 1984, Proc. Natl. Acad. Sci. 81:848-852). The hsp60 and hsp90 families also show similarly high levels of intra-family conservation (Hickey *et al.*, 1989, Mol. Cell. Biol. 9:2615-2626; Jindal, 1989, Mol. Cell. Biol. 9:2279-2283). In addition, it has been discovered that the hsp60, hsp70 and hsp90 families are composed of proteins that are related to the stress proteins in sequence, for  
20 example, having greater than 35% amino acid identity, but whose expression levels are not altered by stress.

Studies of the cellular response to heat shock and other physiological stresses revealed that the hsps are involved not only in cellular protection against these adverse conditions, but also in essential biochemical and immunological processes in unstressed  
25 cells. The hsps accomplish different kinds of chaperoning functions. For example, there has been a suggestion that members of the hsp70 family, located in the cell cytoplasm, nucleus, mitochondria, or endoplasmic reticulum, are involved in the presentation of antigens to the cells of the immune system. Moreover, members of the hsp70 family are known to be involved in the transfer, folding and assembly of proteins in normal cells  
30 (Lindquist *et al.*, 1988, Ann. Rev. Genetics 22:631-677). The hsps are capable of binding proteins or peptides, and of releasing the bound proteins or peptides in the presence of adenosine triphosphate (ATP) or low pH.

#### 2.3.2. Antigen Presentation

Major histocompatibility complex (MHC) molecules present antigens on the  
35 cell surface of antigen-presenting cells. Cytotoxic T lymphocytes (CTLs) then recognize MHC molecules and their associated peptides and kill the target cell. Antigens are

processed by two distinct antigen processing routes depending upon whether their origin is intracellular or extracellular. Intracellular or endogenous protein antigens, *i.e.*, antigens synthesized within the antigen-presenting cell, are presented by MHC class I (MHC I) molecules to CD8+ cytotoxic T lymphocytes. On the other hand, extracellular or  
5 exogenously synthesized antigenic determinants are presented on the cell surface of "specialized" or "professional" APCs (macrophages, for example) by MHC class II molecules to CD4+ T cells (*see generally*, Fundamental Immunology, W.E. Paul (ed.), New York: Raven Press, 1984). This compartmental segregation of antigen processing routes is important to prevent tissue destruction that could otherwise occur during an  
10 immune response as a result of shedding of neighboring cell MHC I antigens.

The heat shock protein gp96 chaperones a wide array of peptides, depending upon the source from which gp96 is isolated (*for review, see* Srivastava *et al.*, 1998, Immunity 8: 657-665). Tumor-derived gp96 carries tumor-antigenic peptides (Ishii *et al.*, 1999, J. Immunology 162:1303-1309), gp96 preparations from virus-infected cells carry  
15 viral epitopes (Suto and Srivastava, 1995, Science 269:1585-1588; Nieland *et al.*, 1998, Proc. Natl. Acad. Sci. USA 95:1800-1805), and gp96 preparations from cells transfected with model antigens such as ovalbumin or  $\beta$ -galactosidase are associated with the corresponding epitopes (Arnold *et al.*, 1995, J. Exp. Med. 182:885-889; Breloer *et al.*, 1998, Eur. J. Immunol. 28:1016-1021). The association of gp96 with peptides occurs *in vivo*  
20 (Menoret and Srivastava, 1999, Biochem. Biophys. Research Commun. 262:813-818). Gp96-peptide complexes, whether isolated from cells (Tamura *et al.*, 1997, Science 278:117-120), or reconstituted *in vitro* (Blachere *et al.*, 1997, J. Exp. Med. 186:1183-1406) are excellent immunogens and have been used extensively to elicit CD8+ T cell responses specific for the gp96-chaperoned antigenic peptides.

25 The capacity of gp96-peptide complexes to elicit an immune response is dependent upon the transfer of the peptide to MHC class I molecules of antigen-presenting cells (Suto and Srivastava, 1995, *supra*). Endogenously synthesized antigens chaperoned by gp96 in the endoplasmic reticulum (ER) can prime antigen-specific CD8+ T cells (or MHC I-restricted CTLs) *in vivo*; this priming of CD8+ T cells requires macrophages.  
30 However, the process whereby exogenously-introduced gp96-peptide complexes elicit the antigen-specific CD8+ T cell response is not completely understood since there is no established pathway for the translocation of extracellular antigens into the class I presentation machinery. Yet antigenic peptides of extracellular origin associated with hsp90 are somehow salvaged by macrophages, channeled into the endogenous pathway, and  
35 presented by MHC I molecules to be recognized by CD8+ lymphocytes (Suto and Srivastava, 1995, *supra*; Blachere *et al.*, 1997, J. Exp. Med. 186:1315-22).

Several models have been proposed to explain the delivery of extracellular peptides for antigen presentation. One proposal, known as the "direct transfer" model, suggests that hsp-chaperoned peptides are transferred to MHC I molecules on the cell surface of macrophages for presentation to CD8<sup>+</sup> T lymphocytes. Another suggestion is that soluble extracellular proteins can be trafficked to the cytosol via constitutive macropinocytosis in bone marrow-derived macrophages and dendritic cells (Norbury *et al.*, 1997, *Eur. J. Immunol.* 27:280-288). Yet another proposed mechanism is that hsps are taken up by the MHC class I molecules of the macrophage, which stimulate the appropriate T cells (Srivastava *et al.*, 1994, *Immunogenetics* 39:93-98). Others have suggested that a novel intracellular trafficking pathway may be involved for the transport of peptides from the extracellular medium into the lumen of the ER (Day *et al.*, 1997, *Proc. Natl. Acad. Sci.* 94:8064-8069; Nicchitta, 1998, *Curr. Opin. in Immunol.* 10:103-109). Further suggestions include the involvement of phagocytes which: (a) possess an ill-defined pathway to shunt protein from the phagosome into the cytosol where it would enter the normal class I pathway; and/or (b) digest ingested material in lysosomes and regurgitate peptides for loading on the surface to class I molecules (Bevan, 1995, *J. Exp. Med.* 182:639-41).

Still others have proposed a receptor-mediated pathway for the delivery of extracellular peptides to the cell surface of APCs for antigen presentation. In view of the extremely small quantity of gp96-chaperoned antigenic peptides required for immunization (Blachere *et al.*, 1997, *supra*), and the strict dependence of immunogenicity of gp96-peptide complexes on functional antigen presenting cells (APCs) (Udono *et al.*, 1994, *Proc. Natl. Acad. Sci. U.S.A.* 91:3077-3081), APCs had been proposed to possess receptors for gp96 (Srivastava *et al.*, 1994, *Immunogenetics* 39:93-98). The mannose receptor is thought to be used in the uptake of gp96 (Ciupitu *et al.*, 1998, *J. Exp. Med.*, 187:685-691). The alpha(2)macroglobulin receptor, also known as CD91, has proven to be a more universal receptor for hsps, with binding to gp96, hsp90, hsp70, and calreticulin having already been demonstrated, as further discussed in Section 2.5 below.

Antigen-presenting cells (APCs), such as macrophages and dendritic cells, are key components of innate and adaptive immune responses. Antigens are generally 'presented' to T cells or B cells on the surfaces of other cells, the APCs. APCs can trap lymph- and blood-borne antigens and, after internalization and degradation, present antigenic peptide fragments, bound to cell-surface molecules of the major histocompatibility complex (MHC), to T cells. APCs may then activate T cells (cell-mediated response) to clonal expansion, and these daughter cells may either develop into cytotoxic T cells or helper T-cells, which in turn activate B (humoral response) cells

with the same MHC-bound antigen to clonal expansion and specific antibody production (see Alberts et al., 1994, Molecular Biology of the Cell, pp. 1238-45).

Stimulation of T cells involves a number of accessory molecules expressed by both T cells and APCs. Co-stimulatory molecules are those accessory molecules that  
5 promote the growth and activation of the T cell, e.g., B7-1, B7-2, CD40, ICAM-1 and MHC II on the APC surface, and CD28, CD40L, T-cell antigen surface receptors (TCRs) and CD4 on the T cell surface (see e.g., Banchereau and Steinman, 1998, Nature 392:245-252). Upon stimulation, co-stimulatory molecules induce release of cytokines, such as interleukin 1 (IL-1) or interleukin 2 (IL-2), interferon, etc., which promote T cell growth and  
10 expression of surface receptors (See e.g., Paul, 1989, Fundamental Immunology, pp. 109-10).

#### 2.4. Adoptive Immunotherapy

Adoptive immunotherapy of cancer refers to a therapeutic approach in which  
15 immune cells with an antitumor reactivity are administered to a tumor-bearing host, with the aim that the cells mediate, either directly or indirectly, the regression of an established tumor. Transfusion of lymphocytes, particularly T lymphocytes, falls into this category and investigators at the National Cancer Institute (NCI) have used autologous reinfusion of peripheral blood lymphocytes or tumor-infiltrating lymphocytes (TIL) T cell cultures from  
20 biopsies of subcutaneous lymph nodules, to treat several human cancers (Rosenberg, S.A., U.S. Patent No. 4,690,914, issued September 1, 1987; Rosenberg, S.A. et al., 1988, *N. England J. Med.* 319:1676-1680). For example, TIL expanded *in vitro* in the presence of interleukin (IL)-2 have been adoptively transferred to cancer patients, resulting in tumor regression in select patients with metastatic melanoma. Melanoma TIL grown in IL-2 have  
25 been identified as activated T-lymphocytes CD3+ HLA-DR+, which are predominantly CD8+ cells with unique *in vitro* antitumor properties. Many long-term melanoma TIL cultures lyse autologous tumors in a specific MHC class I- and T cell antigen receptor dependent manner (Topalian et al., 1989, *J. Immunol.* 142:3714). However, studies of TIL derived from other types of tumors have revealed only scant evidence for cytolytic or  
30 proliferative antitumor immune specificity (Topalian et al., 1990, in *Important Advances in Oncology*, V.T. DeVita, S.A. Hellman and S.A. Rosenberg, eds. J.B. Lippincott, Philadelphia, pp. 19-41). In addition, the toxicity of the high-dose IL-2 + activated lymphocyte treatment advocated by the NCI group has been considerable, including high fevers, severe rigors, hypotension, damage to the endothelial wall due to capillary leak  
35 syndrome, and various adverse cardiac events such as arrhythmias and myocardial infarction (Rosenberg et al., 1988, *N. England J. Med.* 319:1676-1680).



### 2.5. $\alpha$ 2-Macroglobulin

The  $\alpha$ -macroglobulins are members of a protein superfamily of structurally related proteins which also comprises complement components C3, C4 and C5. The human plasma protein alpha(2)macroglobulin ( $\alpha$ 2M) is a 720 kDa homotetrameric protein primarily known as proteinase inhibitor and plasma and inflammatory fluid proteinase scavenger molecule (*for review see* Chu and Pizzo, 1994, Lab. Invest. 71:792). Alpha (2) macroglobulin is synthesized as a 1474 amino acid precursor, the first 23 of which function as a signal sequence that is cleaved to yield a 1451 amino acid mature protein (Kan *et al.*, 1985, Proc. Natl. Acad. Sci. U.S.A. 82:2282-2286).

Alpha(2)macroglobulin promiscuously binds to proteins and peptides with nucleophilic amino acid side chains in a covalent manner (Chu *et al.*, 1994, Ann. N.Y. Acad. Sci. 737:291-307) and targets them to cells which express the  $\alpha$ 2M receptor ( $\alpha$ 2MR) (Chu and Pizzo, 1993, J. Immunol. 150:48). Binding of  $\alpha$ 2M to the  $\alpha$ 2MR is mediated by the C-terminal portion of  $\alpha$ 2M (Holtet *et al.*, 1994, FEBS Lett. 344:242-246) and key residues have been identified (Nielsen *et al.*, 1996, J. Biol. Chem. 271:12909-12912).

Generally known for inhibiting protease activity,  $\alpha$ 2M binds to a variety of proteases thorough multiple binding sites (*see, e.g.*, Hall *et al.*, 1981, Biochem. Biophys. Res. Commun. 100(1):8-16). Protease interaction with  $\alpha$ 2M results in a complex structural rearrangement called transformation, which is the result of a cleavage within the "bait" region of  $\alpha$ 2M after the proteinase becomes "trapped" by thioesters. The conformational change exposes residues required for receptor binding, allowing the  $\alpha$ 2M-proteinase complex to bind to the  $\alpha$ 2MR. Methylamine can induce similar conformational changes and cleavage as that induced by proteinases. The uncleaved form of  $\alpha$ 2M, which is not recognized by the receptor, is often referred to as the "slow" form (s- $\alpha$ 2M). The cleaved form is referred to as the "fast" form (f- $\alpha$ 2M) (reviewed by Chu *et al.*, 1994, Ann. N.Y. Acad. Sci. 737:291-307).

Studies have shown that, in addition to its proteinase-inhibitory functions,  $\alpha$ 2M, when complexed to antigens, can enhance the antigens' ability to be taken up by antigen presenting cells such as macrophages and presented to T cell hybridomas *in vitro* by up to two orders of magnitude (Chu and Pizzo, 1994, Lab. Invest. 71:792), and induce T cell proliferation (Osada *et al.*, 1987, Biochem. Biophys. Res. Commun. 146:26-31). Further evidence suggests that complexing antigen with  $\alpha$ 2M enhances antibody production by crude spleen cells *in vitro* (Osada *et al.*, 1988, Biochem. Biophys. Res. Commun. 150:883) and elicits an *in vivo* antibody responses in experimental rabbits (Chu *et al.*, 1994, J. Immunol. 152:1538-1545) and mice (Mitsuda *et al.*, 1993, Biochem. Biophys. Res.

Commun. 101:1326-1331). However, none of these studies have shown whether  $\alpha$ 2M-antigen complexes are capable of eliciting cytotoxic T cell responses *in vivo*.

$\alpha$ 2M can form complexes with antigens, which are taken up by antigen presenting cells ("APCs") via the  $\alpha$ 2MR, also known as LDL (low-density lipoprotein) Receptor-Related Protein ("LRP") or CD91 (see provisional patent application no. 60/209,266 filed June 2, 2000, which is incorporated by reference herein in its entirety).  $\alpha$ 2M directly competes for the binding of heat shock protein gp96 to the  $\alpha$ 2MR, indicating that  $\alpha$ 2M and hsps may bind to a common recognition site on the  $\alpha$ 2MR (Binder *et al.*, 2000, Nature Immunology 1(2), 151-154). Additionally,  $\alpha$ 2M-antigenic peptide complexes prepared *in vitro* can be administered to animals to generate a cytotoxic T cell response specific to the antigenic molecules (Binder *et al.*, 2001, J. Immunol. 166:4968-72). Thus, because hsps and  $\alpha$ 2M have a number of common functional attributes, such as the ability to bind peptide, the recognition and uptake by the  $\alpha$ 2MR, and the stimulation of a cytotoxic T cell response,  $\alpha$ 2M can be used for immunotherapy against cancer, infectious disease and neurodegenerative or disease.

## 2.6. Saponins

*Quillaja* saponins are a mixture of triterpene glycosides extracted from the bark of the tree *Quillaja saponaria*. They have long been recognized as immune stimulators that can be used as vaccine adjuvants, (Campbell and Peerbaye, 1992, Res. Immunol. 143(5):526-530), and a number of commercially available complex saponin extracts have been utilized as adjuvants. Crude saponins have been extensively employed as adjuvants in veterinary vaccines against foot and mouth disease, and in amplifying the protective immunity conferred by experimental vaccines against protozoal parasites such as *Trypanosoma cruzi* plasmodium and also the humoral response to sheep red blood cells (SRBC) (Bomford, 1982, Int. Arch. Allerg. Appl. Immun. 67:127).

The first commercially available *Quillaja* saponin adjuvants were crude extracts which, because of their variability, were not desirable for use in veterinary practice or in pharmaceutical compositions for man. An early attempt to purify *Quillaja* saponin adjuvants was made by Dalsgaard (1974, Archiv fuer die gesamte Virusforschung 44:243). Dalsgaard partially purified an aqueous extract of the saponin adjuvant material from *Quillaja saponaria* Molina. However, while Dalsgaard's preparation, "Quil-A," was a definite improvement over the previously available commercial saponins, it still exhibited considerable heterogeneity.

Subsequent analysis via high-pressure liquid chromatography showed that Quil A was in fact a heterogeneous mixture of structurally related triterpene glycosides

(U.S. Pat. No. 5,057,540; Kersten *et al.*, 1988, Infect. Immun. 56:432-438; Kensil *et al.*, 1991, J. Immunol. 146:431-437; Kensil *et al.*, 1991, J. Am. Vet. Med. Assoc. 199:1423-1427). However, not all of these saponins were active as adjuvants.

The four most predominant purified *Quillaja* saponins are QS-7, QS-17, 5 QS-18, and QS-21 (alternatively identified as QA-7, QA-17, QA-18, and QA-21). These saponins have been purified by HPLC and low pressure silica chromatography and were found to be adjuvant active, although differing in biological activities such as hemolysis and toxicity in mice. In particular, QS-21 and QS-7 were found to be least toxic in mice (Kensil *et al.*, 1991, J. Immunol. 146:431-437).

10 Due to its potent adjuvant activity and low toxicity, QS-21 (commercially available as the "Stimulon®" adjuvant) has been identified as a useful immunological adjuvant (Kensil *et al.*, 1995, "Structural and Immunological Characterization of the Vaccine Adjuvant QS-21," in Vaccine Design: The Subunit and Adjuvant Approach, Powell and Newman eds., Plenum Press, New York). QS-21 is a complex triterpene glycoside of 15 quillaic acid. QS-21 is glycosylated at triterpene carbon 3, triterpene carbon 28, and carbon 5 of the second fatty acyl unit in a fatty acid domain.

More recently, QS-21 was further purified using hydrophilic interaction chromatography (HILIC) and resolved into two peaks, QS-21-V1 and QS-21-V2, which have been shown to be chemically different compounds. In C57BL/6 mice immunized with 20 vaccines consisting of ovalbumin and either QS-21, QS-1-V1, or QS-21-V2, both of the individual components QS-21-V1 and QS-21-V2 are comparable in adjuvant effect to the original QS-21 peak (containing a mixture of 3:2 QS-21-V1 and QS-21-V2) for boosting the IgG subclasses IgG1, IgG2b, and IgG2 as well as the total IgG titer (U.S. Pat. No. 5,583,112, the entire contents of which are hereby incorporated by reference).

25 *Quillaja* saponins are structurally distinct from the saponins derived from other plant species. Two structural features that distinguish *Quillaja saponaria* saponins from those of other plant species are a fatty acid domain and a triterpene aldehyde at carbon 4 of the triterpene. (Kensil *et al.*, 1995, "Structural and Immunological Characterization of the Vaccine Adjuvant QS-21," in Vaccine Design: The Subunit and Adjuvant Approach, 30 Powell and Newman eds., Plenum Press, New York). Modifications to the aldehyde on the triterpene indicate that this functional group may be involved in the adjuvant mechanism (Soltysik *et al.*, 1995, Vaccine 13(15):1403-1410).

*Quillaja* saponins, particularly QS-7, QS-17, QS-18, and QS-21, have been found to be excellent stimulators of antibody response to soluble T-dependent protein 35 antigens, "subunit antigens", which are poorly immunogenic and require a potent adjuvant for maximization of immune responses. Examples of purified subunit antigens for which

saponin adjuvants that augment the IgG response in mice include keyhole limpet hemocyanin (KLH), HIV-1 gp120 (Bomford *et al.*, 1992, AIDS Res. Hum. Retroviruses 8:1765), and influenza nucleoprotein (Brett *et al.*, 1993, Immunology 80:306). QS-7, QS-17, QS-18 and QS-21 have also been shown to stimulate potent antibody responses in mice to the antigens bovine serum albumin and cytochrome b<sub>5</sub> (Kensil *et al.*, 1991, J. Immunol. 146:431). The level of antibody response induced by these purified saponins was comparable to other commonly used adjuvants, *e.g.*, complete Freund's adjuvant, and superior to aluminum hydroxide.

QS-21 has also been shown to enhance antibody responses to T-independent antigens, including unconjugated bacterial polysaccharides (White *et al.*, 1991, "A purified saponin acts as an adjuvant for a T-independent antigen," in: Immunobiology of Proteins and Peptides, Vol. VI (Atassi ed.), Plenum Press, New York, pp. 207-210). The immunogenicity of the vaccine was further increased by conjugating diphtheria toxoid to the polysaccharide. QS21 enhanced the antibody response to the polysaccharide as well as the carrier, including IgG2a, IgG2b, and IgG3 responses (Coughlin *et al.*, 1995, Vaccine 13(1):17-21).

The ability of adjuvants to modulate the isotype distribution and IgG subclass distribution of antibody response to an antigen through the promotion of Ig subclass switching has important implications for immunity to many bacterial and viral vaccines. QS-7, QS-17, QS-18, and QS-21 stimulate IgG2a response to cytochrome b<sub>5</sub> after administration with saponin doses of 20 µg (Kensil *et al.*, 1991, J. Immunol 146:431). In this regard, QS-21 shifts predominant IgG1 responses to a profile that includes significant IgG2b and IgG2a responses. For example, QS-21 has been shown to stimulate antigen-specific IgG2a to a number of antigens, including *Borrelia burgdorferi* outer surface proteins OspA and OspB (Ma *et al.*, 1994, Vaccine 12(10):925), feline leukemia virus (FeLV) envelope gp70 (Kensil *et al.*, 1991, J. Am. Vet. Med. Assoc. 10:1423), human cytomegalovirus (HCMV) envelope protein gB (Britt *et al.*, 1995, J. Infect. Dis. 171:18), respiratory syncytial virus (RSV) purified fusion protein (Hancock *et al.*, 1995, Vaccine 13(4):391), and tetanus toxoid (Coughlin *et al.*, 1995, Vaccine 13(1):17). QS-21 has also been shown to induce boostable antibody responses (Britt *et al.*, 1995, J. Infect. Dis. 171:18-25; Helling *et al.*, 1995, Cancer Res. 55:2783-2788).

The ability of the QS-21 adjuvant to induce class I major histocompatibility complex (MHC) antigen-restricted cytotoxic T-lymphocyte responses (CTL) after immunization with soluble proteins is a characteristic of saponin adjuvants. A number of studies have shown the ability of QS-21 to induce potent cytotoxic T-lymphocyte (CTL) responses to various antigens, including ovalbumin (Wu *et al.*, 1994, Cell. Immunol.

154:394-406; Newman *et al.*, 1992, J. Immunol. 148(8):2357-2362), recombinant HIV-1 gp160 protein (Wu *et al.*, 1992, J. Immunol. 148:1519), respiratory syncytial virus ("RSV") purified fusion protein (Hancock *et al.*, 1995, Vaccine 13(4):391), and subunit SIV<sub>mac251</sub> gag and env (Newman *et al.*, 1994, AIDS Res. Hum. Retroviruses 10(7):853).

5                   Most of the saponin adjuvant studies have been carried out in mice. However, the adjuvant activity of saponins is not limited to mice; it has also been demonstrated in guinea pigs, rabbits, pigs, sheep, cattle, and nonhuman primates. An adjuvant effect from QS-21 has been observed in cats, guinea pigs, dogs, nonhuman primates, and humans (Kensil *et al.*, 1995, "Structural and Immunological Characterization  
10 of the Vaccine Adjuvant QS-21," in Vaccine Design: The Subunit and Adjuvant Approach, Powell, M. F. and Newman, M. J. eds., Plenum Press, New York).

                  Phase 1 human trials of QS-21 with GM2 ganglioside-keyhole limpet haemocyanin conjugate vaccine have been conducted in patients with malignant melanoma (Livingston *et al.*, 1994, Vaccine 12:1275-1280. Increased immunogenicity after  
15 administration with QS-21 adjuvant was observed (Helling *et al.*, 1995, Cancer Res. 55:2783-2788). In another set of clinical trials, QS-21 was found to be a potent immunological adjuvant that significantly increased the serological response of melanoma patients to the murine antiidiotypic antibody MELIMMUNE-1 (Livingston *et al.*, 1995, Vaccine Res. 4(2):87).

20                   The immune adjuvant effect of saponins is dependent upon dose. Depending upon the antigen and the species, a minimum dose level of QS-21 is required for optimum response (Kensil *et al.*, 1991, J. Immunol.; Kensil *et al.*, 1993, Vaccine Res.; Newman *et al.*, 1992, J. Immunol.; Livingston *et al.*, 1994, Vaccine. Below this minimum dose, the immune adjuvant effect is suboptimal (either low level or absent). QS-7 also has a dose  
25 response curve (Kensil *et al.*, 1991, J. Immunol.).

### 3. Summary of the Invention

                  The present invention provides pharmaceutical compositions comprising an hsp or  $\alpha$ 2M and a saponin, which compositions produce enhanced inhibition of autoimmune  
30 responses when administered in lower doses relative to compositions comprising an hsp or  $\alpha$ 2M in the absence of a saponin. The compositions are therefore useful for treating autoimmune diseases. Accordingly, the present invention provides pharmaceutical compositions comprising a purified hsp and a saponin or a purified  $\alpha$ 2M and a saponin. The pharmaceutical compositions optionally further comprise an antigenic molecule. The  
35 pharmaceutical compositions can be used to treat or prevent an autoimmune disease in an

individual when administered to the individual in an amount effective to treat or prevent the autoimmune disease.

The present invention further provides compositions comprising an hsp or  $\alpha$ 2M, an antigenic molecule, and a saponin, which have enhanced antigenicity or  
5 immunogenicity relative to compositions comprising hsps or  $\alpha$ 2M and antigenic molecules in the absence of a saponin. Such compositions are useful for treating or preventing cancer or an infectious disease or a neurodegenerative or amyloid disease. Accordingly, the present invention provides pharmaceutical compositions comprising a purified hsp or  $\alpha$ 2M, a first antigenic molecule, and a saponin.

10 In one embodiment, the first antigenic molecule is not bound to the saponin or to the hsp or  $\alpha$ 2M. In another embodiment, the pharmaceutical compositions further comprise a second antigenic molecule. The second antigenic molecule can be covalently or non-covalently bound to the hsp or  $\alpha$ 2M. Optionally, the pharmaceutical compositions further comprise a third antigenic molecule, which can be covalently bound to the saponin.

15 In other embodiments, the first antigenic molecule is covalently or non-covalently bound to the hsp or  $\alpha$ 2M. Optionally, the saponin is covalently bound to a second antigenic molecule.

The hsp present in a pharmaceutical composition of the invention can be hsp70, hsp90, gp96, calreticulin, hsp 110, grp170, PDI, or a mixture of two or more of the  
20 foregoing. In certain specific embodiment, the hsp or  $\alpha$ 2M is recombinantly produced as a fusion protein with the second antigenic molecule. In a preferred embodiment, the saponin is QS-7, QS-21, QS-21-V1, or QS-21-V2.

The compositions of the invention comprising hsps or  $\alpha$ 2M, antigenic molecules and saponins can be used to elicit an immune response against cancer or an agent  
25 of infectious disease, by administering to an individual an amount effective to elicit an immune response in the individual. Where eliciting an immune response against a type of cancer is desired, an antigenic molecule is used which displays antigenicity of an antigen of the type of cancer. Where eliciting an immune response against an agent of an infectious disease is desired, an antigenic molecule is used which displays antigenicity of an antigen of  
30 the agent of infectious disease. In other embodiments, the compositions of the invention that comprise an antigenic molecule that displays antigenicity of an antigen of a type of cancer are used to treat or prevent the type of cancer, the compositions of the invention that comprise an antigenic molecule that displays antigenicity of an antigen of an agent of an infectious disease are used to treat or prevent said infectious disease; and the compositions  
35 of the invention that comprise an antigenic molecule associated with a neurodegenerative

disease or an amyloid disease are used to treat or prevent said neurodegenerative or amyloid disease.

Prophylactic and therapeutic dosage regimens and kits are also provided by the invention.

5

#### 4. Detailed Description of the Invention

Methods and pharmaceutical compositions for the prevention and treatment of primary and metastatic neoplastic diseases (also herein referred to as "cancers"), autoimmune diseases, infectious diseases and neurodegenerative and amyloid disease and  
10 for eliciting an immune response in an individual, are described. Particular compositions of the invention and their properties and methods of use are described in the sections and subsections which follow.

The invention is based, in part, on the identification of saponins as useful adjuvants for enhancing the immune response to administration of low doses of hsps or  
15  $\alpha$ 2M alone, or to administration of lower doses of hsps or  $\alpha$ 2M complexed covalently or non-covalently to antigenic molecules. The use of saponins, and in particular QS-7, QS-17, QS-18, QS-21, QS-21-V1, and/or QS-21-V2, permits the use of reduced amounts of hsps or  $\alpha$ 2M and is particularly useful when the quantities of hsps or  $\alpha$ 2M or antigen would otherwise be limiting. The saponins can be used alone or covalently complexed to antigenic  
20 molecules.

The compositions of the invention minimally comprise one or more hsps or  $\alpha$ 2M and a saponin adjuvant. Such compositions, which substantially lack antigenic molecules, are particularly useful in treating an autoimmune disorder. "Antigenic molecule" as used herein refers to a peptide or other molecule with which hsps are  
25 endogenously associated *in vivo* (e.g., in precancerous or cancerous tissue), as well as exogenous antigens/immunogens (i.e., with which the hsps are not complexed *in vivo*) or antigenic/immunogenic fragments and derivatives thereof. Such exogenous antigens and fragments and derivatives (both peptide and non-peptide) thereof for use in complexing with hsps or  $\alpha$ 2M, can be selected from among those known in the art, as well as those  
30 readily identified by standard immunoassays known in the art by detecting the ability to bind antibody or MHC molecules (antigenicity) or generate immune response (immunogenicity). Compositions comprising an hsp or  $\alpha$ 2M and a saponin adjuvant which further comprise an antigenic molecule are useful in eliciting an immune response against the antigenic molecule or a cell in which the antigenic molecule is present, such as a cancer  
35 cell, a cell infected with an infectious organism or a cell or structure, e.g., extracellular deposits or plaques comprising peptide and/or protein fibrils, that displays the hallmarks of

a neurodegenerative or amyloid disease. In certain embodiments, the outcome of eliciting an immune response is prophylaxis or therapy.

The hsps or  $\alpha$ 2M, and/or antigenic molecules are preferably autologous to the individual, although they may also be allogeneic. The hsps or  $\alpha$ 2M and antigenic  
5 molecules can be isolated as naturally-occurring complexes, *e.g.* from cancer cells or cells infected with an infectious agent. Alternatively, the hsps or  $\alpha$ 2M and/or antigenic molecules can be chemically synthesized or recombinantly produced.

Thus, the compositions can be utilized for the prevention of a variety of cancers, *e.g.*, in individuals who are predisposed as a result of familial history or in  
10 individuals with an enhanced risk to cancer due to environmental factors, for the prevention of infectious diseases, *e.g.*, in individuals with enhanced risks of exposure to agents of infectious disease, and for the prevention of neurodegenerative or amyloid diseases, for example in individuals with genetic predispositions to neurodegenerative or amyloid diseases.

15 In certain embodiment, the compositions of the invention are administered in conjunction with administering one or more biological response modifiers, *e.g.*, IFN- $\alpha$ , IFN- $\gamma$ , IL-2, IL-4, IL-6, TNF, or other cytokine growth factors affecting the immune cells.

Optionally, adoptive immunotherapy can be used in conjunction with administration of the hsps or  $\alpha$ 2M, antigenic molecule, and saponin compositions of the  
20 invention in treating or preventing cancer, an infectious disease or neurodegenerative or amyloid disease.

The methods of the invention further comprise methods of making the claimed compositions. The saponin, the antigenic molecule and the hsp or  $\alpha$ 2M are combined under conditions that produce a pharmaceutically effective composition. The  
25 saponin can be combined with the antigenic molecule and then antigen-free hsp or  $\alpha$ 2M is added. Alternatively, the saponin is covalently linked with the antigenic molecule and then combined with antigen-free hsp or  $\alpha$ 2M. In another embodiment, the antigen-free hsp or  $\alpha$ 2M is combined with the antigenic molecule and then the saponin added. In another embodiment, antigen-free hsp or  $\alpha$ 2M, the saponin, and the antigenic molecule are  
30 combined simultaneously. In another embodiment, purified hsp or  $\alpha$ 2M is stripped of bound peptide and antigenic molecule, or antigenic molecule previously covalently linked to saponin, is bound to said hsp or  $\alpha$ 2M *in vitro*. In yet another embodiment, aqueous solutions of the saponin, the antigenic molecule and antigen-free hsp or  $\alpha$ 2M are combined to produce a composition under conditions that permit antigenic molecule binding to the  
35 antigen-free hsp or  $\alpha$ 2M. In yet another embodiment, the saponin is mixed with hsp- or  $\alpha$ 2M-antigenic molecule complexes that were formed endogenously in cells and purified



from the cells. In yet another embodiment, the saponin is mixed with hsp- or  $\alpha$ 2M-antigenic molecule complexes that are produced *in vitro* using purified hsp and antigenic molecule. In certain specific embodiments, the hsp or  $\alpha$ 2M is recombinantly expressed as a fusion protein with an antigenic molecule. In the above methods, multiple hsps (*e.g.* hsp70, hsp90 and gp96),  $\alpha$ 2M, antigenic molecules, and saponins may be combined in a single composition of the invention.

#### 4.1. Compositions of the Invention

The hsp or  $\alpha$ 2M/antigenic molecule/saponin compositions of the invention are administered to elicit an effective specific immune response to the antigenic molecules (and not to the hsp or  $\alpha$ 2M). In accordance with the methods described herein, hsp or  $\alpha$ 2M-antigenic molecule complexes are preferably purified in the range of 60 to 100 percent of the total mg protein, or at least 70%, 80% or 90% of the total mg protein. In another embodiment, hsp or  $\alpha$ 2M-antigenic molecule complexes are purified to apparent homogeneity, as assayed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

In preferred embodiments, non-covalent complexes of hsp70, hsp90, gp96 and/or  $\alpha$ 2M with antigenic molecules are prepared and purified postoperatively from tumor cells obtained from a cancer patient.

In accordance with the methods described herein, immunogenic or antigenic peptides that are endogenously complexed to hsps or  $\alpha$ 2M can be used as specific antigenic molecules. For example, such peptides may be prepared that stimulate cytotoxic T cell responses against different tumor antigens (*e.g.*, tyrosinase, gp100, melan-A, gp75, mucins, *etc.*); viral proteins including, but not limited to, proteins of immunodeficiency virus type I (HIV-I), human immunodeficiency virus type II (HIV-II), hepatitis type A, hepatitis type B, hepatitis type C, influenza, Varicella, adenovirus, herpes simplex type I (HSV-I), herpes simplex type II (HSV-II), rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, papova virus, cytomegalovirus, echinovirus, arbovirus, huntavirus, coxsackie virus, mumps virus, measles virus, rubella virus and polio virus; antigenic molecules of neurodegenerative or amyloid diseases, such as  $\beta$ -amyloid or a fragment thereof, an oligomeric A $\beta$  complex or a fragment thereof, an ApoE4-A $\beta$  complex or a fragment thereof, tau protein or a fragment thereof, a mutant amyloid precursor protein or a fragment thereof, a mutant of presenillin or a fragment thereof,  $\alpha$ -synuclein or a fragment thereof, or a prion protein, and their antigenic derivatives. In the embodiment wherein the antigenic molecules are peptides noncovalently complexed to hsps or  $\alpha$ 2M *in vivo*, the complexes can be isolated from cells, or alternatively, produced *in vitro* from purified preparations each of hsps or  $\alpha$ 2M and antigenic molecules.

In another specific embodiment, specific antigens of cancers (*e.g.*, tumors) or infectious agents (*e.g.*, viral antigens, bacterial antigens, etc.) can be obtained by purification from natural sources, by chemical synthesis, or recombinantly, and, through *in vitro* procedures such as that described below, noncovalently complexed to hsps or  $\alpha 2M$ .

- 5 In another embodiment, the hsp or  $\alpha 2M$  -specific antigen complexes or antigens obtained by purification from natural sources, by chemical synthesis, or recombinantly, and, through *in vitro* procedures, are related to an autoimmune disease. These autoimmune diseases include, but are not limited to, insulin-dependent diabetes mellitus (*i.e.*, IDDM, or autoimmune diabetes), multiple sclerosis, systemic lupus
- 10 erythematosus, Sjogren's syndrome, scleroderma, polymyositis, chronic active hepatitis, mixed connective tissue disease, primary biliary cirrhosis, pernicious anemia, autoimmune thyroiditis, idiopathic Addison's disease, vitiligo, gluten-sensitive enteropathy, Graves' disease, myasthenia gravis, autoimmune neutropenia, idiopathic thrombocytopenia purpura, rheumatoid arthritis, cirrhosis, pemphigus vulgaris, autoimmune infertility, Goodpasture's
- 15 disease, bullous pemphigoid, discoid lupus, ulcerative colitis, and dense deposit disease. The diseases set forth above, as referred to herein, include those exhibited by animal models for such diseases, such as, for example non-obese diabetic (NOD) mice for IDDM and experimental autoimmune encephalomyelitis (EAE) mice for multiple sclerosis.

- In an embodiment wherein an  $\alpha 2M$  or hsp -specific antigenic molecule
- 20 complex to be used is a complex that is produced *in vivo* in cells, exemplary purification procedures such as described in Sections 4.2.1 and 4.2.2 below can be employed. Alternatively, in an embodiment wherein one wishes to use specific antigenic molecules by complexing to hsps *in vitro*, hsps can be purified for such use from the endogenous hsp-peptide complexes in the presence of ATP or low pH (or chemically synthesized or
- 25 recombinantly produced). The protocols described herein may be used to isolate hsp-peptide complexes, or the hsps alone, from any eukaryotic cells for example, tissues, isolated cells, or immortalized eukaryote cell lines infected with a preselected intracellular pathogen, tumor cells or tumor cell lines.

- In another embodiment, part or all of the first antigen is covalently bound to
- 30 an hsp to form an hsp-first antigen complex by any means known in the art. In a preferred embodiment, the amount of hsp-first antigen complex is about 0.1  $\mu g$  or greater. In an alternative first embodiment,  $\alpha 2M$  is used in place of hsp. In a preferred alternative embodiment, the hsp or  $\alpha 2M$  and specific antigen are expressed as a recombinant fusion protein. To produce such a recombinant fusion protein, an expression vector is constructed
- 35 using nucleic acid sequences encoding  $\alpha 2M$  fused to sequences encoding an antigenic molecule, using recombinant methods known in the art (*see* Suzue *et al.*, 1997, Proc. Natl.

Acad. Sci. U.S.A. 94: 13146-51).  $\alpha$ 2M-antigenic peptide fusions are then expressed and isolated. By specifically designing the antigenic peptide portion of the molecule, such fusion proteins can be used to elicit an immune response and in immunotherapy against target cancer and infectious diseases.

5           In another embodiment, part or all of the first antigen is noncovalently bound to an hsp to form an hsp-first antigen complex. In a preferred embodiment, the amount of hsp-first antigen complex is about 0.1  $\mu$ g or greater. In an alternative second embodiment,  $\alpha$ 2M is used in place of hsp.

10           In another embodiment, the part or all of the first antigen is covalently bound to a saponin.

          In another embodiment, the pharmaceutical compositions of the invention for the prevention or treatment of an autoimmune disorder comprise an hsp or  $\alpha$ 2M and may further comprise an antigen that may not elicit a specific immune response to an autoimmune disease.

15           In another embodiment, the pharmaceutical compositions of the previous embodiments may include unbound antigen (*i.e.* antigen not complexed with an hsp or  $\alpha$ 2M).

          In one embodiment, the amount of saponin in a pharmaceutical composition of the invention is about 1 micrograms or more. In a preferred embodiment, the amount of  
20 saponin in the pharmaceutical composition is from about 10 to about 1000 micrograms. In a particularly preferred embodiment, the amount of saponin is about 100 micrograms.

          In other embodiments, the saponin is QS-7, QS-21, QS-21-V1, or QS-21-V2. In a preferred embodiment, the saponin is QS-21. The amount of QS-21 in the pharmaceutical compositions is preferably about 1 microgram or more. In a particularly  
25 preferred embodiment, the amount of QS-21 is from about 10 to about 1000 micrograms. In a particularly preferred embodiment, the amount of QS-21 is about 100 micrograms.

          In another embodiment, the pharmaceutical composition comprises an hsp-first antigen complex wherein the first antigen is either covalently or noncovalently bound to the hsp, an hsp-second antigen complex wherein the second antigen is either covalently  
30 or noncovalently bound to the hsp, and a saponin selected from the group consisting of QS-7, QS-21, QS-21-V1, and QS-21-V2, wherein the amount of specific hsp-first antigen complex is about 0.1  $\mu$ g or greater and the amount of QS-21 is about 1 microgram or greater.

          In another embodiment, the pharmaceutical composition comprises an  $\alpha$ 2M-  
35 first antigen complex wherein the first antigen is either covalently or noncovalently bound to the  $\alpha$ 2M and QS-21, wherein the amount of QS-21 is about 1 microgram or greater.

In a particularly preferred form of the above hsp-containing embodiments, the hsp is selected from the group consisting of hsp70, hsp90, gp96, calreticulin, hsp 110, grp170, PDI, or a mixture of two or more of the foregoing hsps.

In another particularly preferred form of the above embodiments, an hsp or  
5  $\alpha$ 2M is purified from cancerous or infected tissues.

In another particularly preferred form of the above embodiments, an hsp or  $\alpha$ 2M is purified from a cell line. In a specific embodiment, the cell line is transfected with a nucleic acid encoding the hsp or  $\alpha$ 2M. In other specific embodiments, the cell line is transfected with a nucleic acid encoding the antigenic molecule and may further be  
10 transfected with a nucleic acid encoding the hsp or  $\alpha$ 2M.

In another particularly preferred form of the above embodiments, an hsp or  $\alpha$ 2M is a recombinant protein.

In another particularly preferred form of the above embodiments, the first and/or second antigen, when present in the composition, is a synthetic or recombinantly  
15 generated peptide.

In another particularly preferred form of the above embodiments, the first antigen and/or second antigen, when present in the composition, is bound to an hsp or  $\alpha$ 2M *in vitro*.

The present invention also encompasses methods for making pharmaceutical  
20 compositions comprising (a) an hsp or  $\alpha$ 2M, (b) a first antigen that will elicit a specific immune response to a cancer, infectious disease, neurodegenerative or amyloid disease, or autoimmune disease, and (c) a saponin adjuvant, said methods comprising combining the saponin, the first antigen and the hsp or  $\alpha$ 2M under conditions that produce a pharmaceutically effective composition.

In one embodiment, the saponin is combined with a specific first antigen and then with antigen-free hsp or  $\alpha$ 2M.

In another embodiment, the saponin is covalently linked with a specific first antigen and then combined with antigen-free hsp or  $\alpha$ 2M (*i.e.* stripped of endogenous antigenic molecule). In a preferred embodiment, the antigen covalently linked to the  
30 saponin is combined with the antigen-free hsp or  $\alpha$ 2M under conditions that allow binding of antigen to hsp or  $\alpha$ 2M.

In another embodiment, antigen-free hsp or  $\alpha$ 2M is combined with a specific first antigen and then with the saponin. In a preferred embodiment, the specific first antigen is combined with the antigen-free hsp or  $\alpha$ 2M under conditions that allow binding of  
35 antigen to hsp.

In another embodiment, antigen-free hsp or  $\alpha$ 2M, a saponin, and a specific first antigen are combined simultaneously.

Another embodiment comprises combining aqueous solutions of saponin, antigen, and antigen-free hsp or  $\alpha$ 2M, under conditions that permit antigen binding to the antigen-free hsp or  $\alpha$ 2M.

In the compositions of the invention, the presence of saponins does not adversely affect the ability of hsp or  $\alpha$ 2M to bind or retain noncovalently-bound antigenic molecules.

#### 4.2. Purification of $\alpha$ 2M and hsp Complexes

The following  $\alpha$ 2M or hsp polypeptides stripped of antigenic molecules, and  $\alpha$ 2M-antigenic molecule complexes or hsp-antigenic molecule complexes, can be used together with one or more saponins in the compositions of the invention.

##### 4.2.1. $\alpha$ 2M Antigenic Molecule Complexes

Described below are methods for purifying  $\alpha$ 2M polypeptides or  $\alpha$ 2M polypeptide-antigenic molecule complexes for use in the invention from recombinant cells, and, with minor modifications known in the art, the  $\alpha$ 2M polypeptide or  $\alpha$ 2M-antigenic molecule complexes from cell culture. Recombinant cells include, for example, cells expressing antigenic molecules and recombinantly expressing an  $\alpha$ 2M polypeptide. Such cells may be derived from a variety of sources, including, but not limited to, cells infected with an infectious agent and cancer cells.

Methods are disclosed for purification of recombinant  $\alpha$ 2M polypeptide-antigenic molecule complexes by affinity purification, based on the properties of the affinity label present on the  $\alpha$ 2M polypeptide. One approach is based on specific molecular interactions between a tag and its binding partner. The other approach relies on the immunospecific binding of an antibody to an epitope present on the tag. The principle of affinity chromatography well known in the art is generally applicable to both of these approaches.

To produce  $\alpha$ 2M polypeptide-antigenic molecule complexes, a nucleotide sequence encoding an  $\alpha$ 2M polypeptide can be introduced into a cell. When an antigenic molecule is present in the cell, the  $\alpha$ 2M polypeptide can associate intracellularly with the antigenic molecule, forming a covalent or a non-covalent complex of  $\alpha$ 2M polypeptide and the antigenic molecule. Cells into which an  $\alpha$ 2M polypeptide-encoding nucleotide sequence can be introduced, include, but are not limited to, epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes,

B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.*, as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, *etc.* The choice of cell type depends on the type of tumor, infectious  
5 disease or neurodegenerative or amyloid disease being treated or prevented, and can be determined by one of skill in the art.

For example, an expression construct comprising a nucleic acid sequence encoding the  $\alpha 2M$  polypeptide is introduced into an antigenic cell. As used herein, antigenic cells may include cells that are infected with an infectious agent or pathogen, cells  
10 infected with non-infectious or non-pathogenic forms of an infectious agent or pathogen (*e.g.*, by use of a helper infectious agent), tissue or cell culture model systems for neurodegenerative or amyloid diseases, cells infected by or engineered to express an attenuated form of an infectious agent or a non-pathogenic or replication-deficient variant of a pathogen, pre-neoplastic cells that are infected with a cancer-causing infectious agent,  
15 such as a virus, but which are not yet neoplastic; or antigenic cells that have been exposed to a mutagen or cancer-causing agent, such as, for example DNA-damaging agents, radiation, *etc.* Other cells that can be used are pre-neoplastic cells which are in transition from a normal to a neoplastic form as characterized by morphology, physiological or biochemical functions. Preferably, the cancer cells and pre-neoplastic cells used in the  
20 methods of the invention are of mammalian origin. Mammals contemplated by this aspect of the invention include humans, companion animals (*e.g.*, dogs and cats), livestock animals (*e.g.*, sheep, cattle, goats, pigs and horses), laboratory animals (*e.g.*, mice, rats and rabbits), and captive or free wild animals.

Virtually any cancer cell, preferably a human cancer cell, can be used in the  
25 present methods for producing  $\alpha 2M$  polypeptide-antigenic molecule complexes. The cancer cells provide the antigenic peptides which become associated covalently or noncovalently with the expressed  $\alpha 2M$  polypeptide.  $\alpha 2M$  polypeptide-antigenic molecule complexes are then purified from the cells and used to treat such cancers. Cancers which can be treated or prevented with immunogenic compositions prepared by methods of the  
30 invention include, but are not limited to, tumors such as sarcomas and carcinomas. Examples of cancers that are amenable to the methods of the invention are listed in the Target Cancers Section below. Accordingly, any tissues or cells isolated from a pre-neoplastic lesion, a cancer, including cancer that has metastasized to multiple remote sites, can be used in the present method. For example, cells found in abnormally growing tissue,  
35 circulating leukemic cells, metastatic lesions as well as solid tumor tissue can be used.

Cell lines derived from a pre-neoplastic lesion, cancer tissues or cancer cells can also be used, provided that the cells of the cell line have at least one or more antigenic determinants in common with antigens on the target cancer cells. Cancer tissues, cancer cells, cells infected with a cancer-causing agent, other pre-neoplastic cells, and cell lines of human origin are preferred.

Cancer and pre-neoplastic cells can be identified by any method known in the art. For example, cancer cells can be identified by morphology, enzyme assays, proliferation assays, cytogenetic characterization, DNA mapping, DNA sequencing, the presence of cancer-causing virus, or a history of exposure to mutagen or cancer-causing agent, imaging, *etc.* Cancer cells may also be obtained by surgery, endoscopy, or other biopsy techniques. If some distinctive characteristics of the cancer cells are known, they can also be obtained or purified by any biochemical or immunological methods known in the art, such as but not limited to affinity chromatography, and fluorescence activated cell sorting (*e.g.*, with fluorescently tagged antibody against an antigen expressed by the cancer cells).

Cancer tissues, cancer cells or cell lines may be obtained from a single individual or pooled from several individuals. It is not essential that clonal, homogeneous, or purified population of cancer cells be used. It is also not necessary to use cells of the ultimate target *in vivo* (*e.g.*, cells from the tumor of the intended recipient), so long as at least one or more antigenic determinants on the target cancer cells is present on the cells used for expression of the  $\alpha 2M$  polypeptide. In addition, cells derived from distant metastases may be used to prepare an immunogenic composition against the primary cancer. A mixture of cells can be used provided that a substantial number of cells in the mixture are cancer cells and share at least one antigenic determinant with the target cancer cell. In a specific embodiment, the cancer cells to be used in expressing an  $\alpha 2M$  polypeptide are purified.

#### 4.2.1.1 *In Vitro* Complexing

Complexes of  $\alpha 2M$  polypeptides and antigenic molecules may be produced *in vitro*. Immunogenic  $\alpha 2M$  polypeptide-antigenic molecule complexes can be generated *in vitro* by coupling of an  $\alpha 2M$  polypeptide with an antigenic peptide. Procedures for forming such  $\alpha 2M$ -antigenic molecule complexes and methods for isolating antigenic peptides are described below.

In general, when an  $\alpha 2M$  is mixed with a protease, cleavage of the "bait" region of  $\alpha 2M$  takes place, the proteinase becomes "trapped" by thioesters; and a conformational change takes place that allows binding of the  $\alpha 2M$  complex to the  $\alpha 2M$

receptor. During proteolytic activation of  $\alpha 2M$ , non-proteolytic ligands can become covalently bound to the activated thioesters. Non-proteolytic ligands can also be incorporated into the activated  $\alpha 2M$  molecule by ammonia or methylamine during reversal of the nucleophilic activation, employing heat (Grøn and Pizzo, 1998, Biochemistry, 37: 6009-6014). Such conditions that allow fortuitous trapping of peptides by  $\alpha 2M$  are employed to prepare the  $\alpha 2M$  -antigenic complexes for use in the invention. Methods for such covalent coupling have been described previously (Osada *et al.*, 1987, Biochem. Biophys. Res. Commun. 146:26-31; Osada *et al.*, 1988, Biochem. Biophys. Res. Commun. 150:883; Chu and Pizzo, 1993, J. Immunol. 150:48; Chu *et al.*, 1994, Ann. N.Y. Acad. Sci. 737:291-307; Mitsuda *et al.*, 1993, Biochem. Biophys. Res. Commun. 101:1326-1331). Thus in one embodiment, an  $\alpha 2M$  antigenic molecule complex can be prepared as described by Grøn and Pizzo, 1998, Biochemistry, 37: 6009-6014. The method of Grøn and Pizzo yields complexes of  $\alpha 2M$  that are covalently bound to antigenic molecules.

For example,  $\alpha 2M$  polypeptide is mixed with an antigenic molecule in the presence of a protease, ammonia or other small amine nucleophiles such as methylamine and ethylamine. Non-limiting examples of proteases which may be used include trypsin, porcine pancreatic elastase (PEP), human neutrophil elastase, cathepsin G, *S. aureus* V-8 proteinase trypsin,  $\alpha$ -chymotrypsin, V8 protease, papain, and proteinase K (see Ausubel *et al.*, eds., in "Current Protocols in Molecular Biology", Greene Publishing Associates and Wiley Interscience, New York, 17.4.6-17.4.8). A preferred, exemplary protocol for complexing an  $\alpha 2M$  polypeptide and an antigenic molecule *in vitro* follows. The antigenic molecules (1  $\mu$ g -20 mg) and the  $\alpha 2M$  polypeptide (1  $\mu$ g-20 mg) are mixed together in phosphate-buffered saline (PBS) (100  $\mu$ l - 5 ml) in the presence of a protease, such as trypsin (0.92 mg trypsin in approximately 500  $\mu$ l PBS, to give an approximately 5:1 antigenic molecule :  $\alpha 2M$  polypeptide molar ratio. The mixture is then incubated for 5-15 minutes at 37°C. 500  $\mu$ l 4 mg/ml p-Aphenyl methyl sulfonyl fluoride (p-APMSF) is added to the solution to inhibit trypsin activity and incubated for 2 hrs at 25°C. The preparations can be centrifuged through a Centricon 10 assembly (Millipore) to remove any unbound peptide. Alternatively, free antigenic molecule may be removed by passage over a gel permeation column. The association of the peptides with the  $\alpha 2M$  polypeptide can be assayed by SDS-PAGE. This is the preferred method for *in vitro* complexing of antigenic molecules isolated from MHC-antigenic molecule complexes, or peptides disassociated from endogenous  $\alpha 2M$ -antigenic molecule complexes.

In a more preferred method, an  $\alpha 2M$ -antigenic molecule complex is prepared according to the method described by Blachere *et al.*, J. Exp. Med. 186(8):1315-22, which incorporated by reference hereinf in its entirety for *in vitro* complexing of hsp's to antigenic



molecule, with the hsp component substituted by  $\alpha 2M$ . The Blachere method yields complexes of  $\alpha 2M$  bound to antigenic molecules.

Antigenic molecules may be isolated from various sources, chemically synthesized, or produced recombinantly. Such methods can be readily adapted for medium  
5 or large scale production of the immunotherapeutic or prophylactic vaccines.

Following complexing, the immunogenic  $\alpha 2M$ -antigenic molecule complexes can optionally be assayed *in vitro* using, for example, the mixed lymphocyte target cell assay (MLTC) described below. Once immunogenic complexes have been isolated they can be optionally characterized further in animal models using the preferred  
10 administration protocols and excipients discussed below.

#### 4.2.2. Preparation and Purification of hsp70-peptide Complexes

The purification of hsp70-peptide complexes has been described previously,  
15 see, for example, Udoni et al., 1993, *J. Exp. Med.* 178:1391-1396. A procedure that may be used, presented by way of example but not limitation, is as follows:

Initially, tumor cells are suspended in 3 volumes of 1X Lysis buffer consisting of 5mM sodium phosphate buffer, pH 7, 150mM NaCl, 2mM  $CaCl_2$ , 2mM  $MgCl_2$  and 1mM phenyl methyl sulfonyl fluoride (PMSF). Then, the pellet is sonicated, on  
20 ice, until >99% cells are lysed as determined by microscopic examination. As an alternative to sonication, the cells may be lysed by mechanical shearing and in this approach the cells typically are resuspended in 30mM sodium bicarbonate pH 7.5, 1mM PMSF, incubated on ice for 20 minutes and then homogenized in a Dounce homogenizer until >95% cells are lysed.

Then the lysate is centrifuged at 1,000g for 10 minutes to remove unbroken  
25 cells, nuclei and other cellular debris. The resulting supernatant is recentrifuged at 100,000g for 90 minutes, the supernatant harvested and then mixed with Con A Sepharose equilibrated with phosphate buffered saline (PBS) containing 2mM  $Ca^{2+}$  and 2mM  $Mg^{2+}$ . When the cells are lysed by mechanical shearing the supernatant is diluted with an equal  
30 volume of 2X lysis buffer prior to mixing with Con A Sepharose. The supernatant is then allowed to bind to the Con A Sepharose for 2-3 hours at 4°C. The material that fails to bind is harvested and dialyzed for 36 hours (three times, 100 volumes each time) against 10mM Tris-Acetate pH 7.5, 0.1mM EDTA, 10mM NaCl, 1mM PMSF. Then the dialyzate is centrifuged at 17,000 rpm (e.g. Sorvall SS34 rotor) for 20 minutes. Then the resulting  
35 supernatant is harvested and applied to a Mono Q FPLC column equilibrated in 20mM Tris-Acetate pH 7.5, 20mM NaCl, 0.1mM EDTA and 15mM 2-mercaptoethanol. The column is then developed with a 20mM to 500mM NaCl gradient and then eluted fractions

fractionated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and characterized by immunoblotting using an appropriate anti-hsp70 antibody (such as from clone N27F3-4, from StressGen).

Fractions strongly immunoreactive with the anti-hsp70 antibody are pooled  
5 and the hsp70-peptide complexes precipitated with ammonium sulfate; specifically with a 50%-70% ammonium sulfate cut. The resulting precipitate is then harvested by centrifugation at 17,000 rpm (SS34 Sorvall rotor) and washed with 70% ammonium sulfate. The washed precipitate is then solubilized and any residual ammonium sulfate removed by gel filtration on a Sephadex<sup>R</sup> G25 column (Pharmacia). If necessary the hsp70 preparation  
10 thus obtained can be repurified through the Mono Q FPLC Column as described above.

The hsp70-peptide complex can be purified to apparent homogeneity using this method. Typically 1 mg of hsp70-peptide complex can be purified from 1 g of cells/tissue.

An improved method for purification of hsp70-peptide complexes comprises  
15 contacting cellular proteins with ADP or a nonhydrolyzable analog of ATP affixed to a solid substrate, such that hsp70 in the lysate can bind to the ADP or nonhydrolyzable ATP analog, and eluting the bound hsp70. A preferred method uses column chromatography with ADP affixed to a solid substratum (*e.g.*, ADP-agarose). The resulting hsp70  
preparations are higher in purity and devoid of contaminating peptides. The hsp70 yields  
20 are also increased significantly by about more than 10 fold. Alternatively, chromatography with nonhydrolyzable analogs of ATP, instead of ADP, can be used for purification of hsp70-peptide complexes. By way of example but not limitation, purification of hsp70-peptide complexes by ADP-agarose chromatography can be carried out as follows:

Meth A sarcoma cells (500 million cells) are homogenized in hypotonic  
25 buffer and the lysate is centrifuged at 100,000 g for 90 minutes at 4°C. The supernatant is applied to an ADP-agarose column. The column is washed in buffer and is eluted with 5 column volumes of 3 mM ADP. The hsp70-peptide complexes elute in fractions 2 through 10 of the total 15 fractions which elute. The eluted fractions are analyzed by SDS-PAGE. The hsp70-peptide complexes can be purified to apparent homogeneity using this  
30 procedure.

#### 4.2.3. Preparation and Purification of hsp90-peptide Complexes

A procedure that can be used, presented by way of example and not  
35 limitation, is as follows:

Initially, tumor cells are suspended in 3 volumes of 1X Lysis buffer consisting of 5mM sodium phosphate buffer (pH7), 150mM NaCl, 2mM CaCl<sub>2</sub>, 2mM

MgCl<sub>2</sub> and 1mM phenyl methyl sulfonyl fluoride (PMSF). Then, the pellet is sonicated, on ice, until >99% cells are lysed as determined by microscopic examination. As an alternative to sonication, the cells may be lysed by mechanical shearing and in this approach the cells typically are resuspended in 30mM sodium bicarbonate pH 7.5, 1mM PMSF, incubated on ice for 20 minutes and then homogenized in a Dounce homogenizer until >95% cells are lysed.

Then the lysate is centrifuged at 1,000g for 10 minutes to remove unbroken cells, nuclei and other cellular debris. The resulting supernatant is recentrifuged at 100,000g for 90 minutes, the supernatant harvested and then mixed with Con A Sepharose equilibrated with PBS containing 2mM Ca<sup>2+</sup> and 2mM Mg<sup>2+</sup>. When the cells are lysed by mechanical shearing the supernatant is diluted with an equal volume of 2X Lysis buffer prior to mixing with Con A Sepharose. The supernatant is then allowed to bind to the Con A Sepharose for 2-3 hours at 4°C. The material that fails to bind is harvested and dialyzed for 36 hours (three times, 100 volumes each time) against 10mM Tris-Acetate pH 7.5, 0.1mM EDTA, 10mM NaCl, 1mM PMSF. Then the dialyzate is centrifuged at 17,000 rpm (Sorvall SS34 rotor) for 20 minutes. Then the resulting supernatant is harvested and applied to a Mono Q FPLC column equilibrated with lysis buffer. The proteins are then eluted with a salt gradient of 200mM to 600mM NaCl.

The eluted fractions are fractionated by SDS-PAGE and fractions containing the hsp90-peptide complexes identified by immunoblotting using an anti-hsp90 antibody such as 3G3 (Affinity Bioreagents). hsp90-peptide complexes can be purified to apparent homogeneity using this procedure. Typically, 150-200 µg of hsp90-peptide complex can be purified from 1g of cells/tissue.

#### 4.2.4. Preparation and Purification of gp96-peptide Complexes

A procedure that can be used, presented by way of example and not limitation, is as follows:

A pellet of tumors is resuspended in 3 volumes of buffer consisting of 30mM sodium bicarbonate buffer (pH 7.5) and 1mM PMSF and the cells allowed to swell on ice 20 minutes. The cell pellet is then homogenized in a Dounce homogenizer (the appropriate clearance of the homogenizer will vary according to each cell type) on ice until >95% cells are lysed.

The lysate is centrifuged at 1,000g for 10 minutes to remove unbroken cells, nuclei and other debris. The supernatant from this centrifugation step is then recentrifuged at 100,000g for 90 minutes. The gp96-peptide complex can be purified either from the 100,000 pellet or from the supernatant.

When purified from the supernatant, the supernatant is diluted with equal volume of 2X lysis buffer and the supernatant mixed for 2-3 hours at 4°C with Con A Sepharose equilibrated with PBS containing 2mM  $\text{Ca}^{2+}$  and 2mM  $\text{Mg}^{2+}$ . Then, the slurry is packed into a column and washed with 1X lysis buffer until the  $\text{OD}_{280}$  drops to baseline.

- 5 Then, the column is washed with 1/3 column bed volume of 10%  $\alpha$ -methyl mannoside ( $\alpha$ -MM) dissolved in PBS containing 2mM  $\text{Ca}^{2+}$  and 2mM  $\text{Mg}^{2+}$ , the column sealed with a piece of parafilm, and incubated at 37°C for 15 minutes. Then the column is cooled to room temperature and the parafilm removed from the bottom of the column. Five column volumes of the  $\alpha$ -MM buffer are applied to the column and the eluate analyzed by SDS-  
10 PAGE. Typically the resulting material is about 60-95% pure, however this depends upon the cell type and the tissue-to-lysis buffer ratio used. Then the sample is applied to a Mono Q FPLC column (Pharmacia) equilibrated with a buffer containing 5mM sodium phosphate, pH 7. The proteins are then eluted from the column with a 0-1M NaCl gradient and the gp96 fraction elutes between 400mM and 550mM NaCl.

- 15 The procedure, however, may be modified by two additional steps, used either alone or in combination, to consistently produce apparently homogeneous gp96-peptide complexes. One optional step involves an ammonium sulfate precipitation prior to the Con A purification step and the other optional step involves DEAE-Sepharose purification after the Con A purification step but before the Mono Q FPLC step.

- 20 In the first optional step, described by way of example as follows, the supernatant resulting from the 100,000g centrifugation step is brought to a final concentration of 50% ammonium sulfate by the addition of ammonium sulfate. The ammonium sulfate is added slowly while gently stirring the solution in a beaker placed in a tray of ice water. The solution is stirred from about 1/2 to 12 hours at 4°C and the resulting  
25 solution centrifuged at 6,000 rpm (Sorvall SS34 rotor). The supernatant resulting from this step is removed, brought to 70% ammonium sulfate saturation by the addition of ammonium sulfate solution, and centrifuged at 6,000 rpm (Sorvall SS34 rotor). The resulting pellet from this step is harvested and suspended in PBS containing 70% ammonium sulfate in order to rinse the pellet. This mixture is centrifuged at 6,000 rpm  
30 (Sorvall SS34 rotor) and the pellet dissolved in PBS containing 2mM  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ . Undissolved material is removed by a brief centrifugation at 15,000 rpm (Sorvall SS34 rotor). Then, the solution is mixed with Con A Sepharose and the procedure followed as before.

- In the second optional step, described by way of example as follows, the  
35 gp96 containing fractions eluted from the Con A column are pooled and the buffer exchanged for 5mM sodium phosphate buffer, pH 7, 300mM NaCl by dialysis, or

preferably by buffer exchange on a Sephadex G25 column. After buffer exchange, the solution is mixed with DEAE-Sepharose previously equilibrated with 5mM sodium phosphate buffer, pH 7, 300mM NaCl. The protein solution and the beads are mixed gently for 1 hour and poured into a column. Then, the column is washed with 5mM sodium phosphate buffer, pH 7, 300mM NaCl, until the absorbance at 280nm drops to baseline. Then, the bound protein is eluted from the column with five volumes of 5mM sodium phosphate buffer, pH 7, 700mM NaCl. Protein containing fractions are pooled and diluted with 5mM sodium phosphate buffer, pH 7 in order to lower the salt concentration to 175mM. The resulting material then is applied to the Mono Q FPLC column (Pharmacia) equilibrated with 5mM sodium phosphate buffer, pH 7 and the protein that binds to the Mono Q FPLC column (Pharmacia) is eluted as described before.

It is appreciated, however, that one skilled in the art may assess, by routine experimentation, the benefit of incorporating the second optional step into the purification protocol. In addition, it is appreciated also that the benefit of adding each of the optional steps will depend upon the source of the starting material.

When the gp96 fraction is isolated from the 100,000g pellet, the pellet is suspended in 5 volumes of PBS containing either 1% sodium deoxycholate or 1% octyl glucopyranoside (but without the  $Mg^{2+}$  and  $Ca^{2+}$ ) and incubated on ice for 1 hour. The suspension is centrifuged at 20,000g for 30 minutes and the resulting supernatant dialyzed against several changes of PBS (also without the  $Mg^{2+}$  and  $Ca^{2+}$ ) to remove the detergent. The dialysate is centrifuged at 100,000g for 90 minutes, the supernatant harvested, and calcium and magnesium are added to the supernatant to give final concentrations of 2mM, respectively. Then the sample is purified by either the unmodified or the modified method for isolating gp96-peptide complex from the 100,000g supernatant, see above.

The gp96-peptide complexes can be purified to apparent homogeneity using this procedure. About 10-20µg of gp96 can be isolated from 1g cells/tissue.

#### **4.2.5. Preparation and Purification of hsp110-peptide Complexes**

A procedure, described by Wang *et al.*, 2001, J. Immunol. 166(1):490-7, that can be used, presented by way of example and not limitation, is as follows:

A pellet (40-60 ml) of cell or tissue, *e.g.*, tumor cell tissue, is homogenized in 5 vol of hypotonic buffer (30 mM sodium bicarbonate, pH7.2, and protease inhibitors) by Dounce homogenization. The lysate is centrifuged at  $4,500 \times g$  and then  $100,000 \times g$  for 2 hours. If the cells or tissues are of hepatic origin, the resulting supernatant is first applied to a blue Sepharose column (Pharmacia) to remove albumin. Otherwise, the resulting supernatant is applied to a Con A-Sepharose column (Pharmacia Biotech,

Piscataway, NJ) previously equilibrated with binding buffer (20mM Tris-HCl, pH 7.5; 100mM NaCl; 1mM MgCl<sub>2</sub>; 1 mM CaCl<sub>2</sub>; 1 mM MnCl<sub>2</sub>; and 15 mM 2-ME). The bound proteins are eluted with binding buffer containing 15%  $\alpha$ -D-o-methylmannoside (Sigma, St. Louis, MO).

5                   Con A-Sepharose unbound material is first dialyzed against a solution of 20 mM Tris-HCl, pH 7.5; 100 mM NaCl; and 15 mM 2-ME, and then applied to a DEAE-Sepharose column and eluted by salt gradient from 100 to 500 mM NaCl. Fractions containing hsp110 are collected, dialyzed, and loaded onto a Mono Q (Pharmacia) 10/10 column equilibrated with 20mM Tris-HCl, pH 7.5; 200 mM NaCl; and 15 mM 2-ME. The  
10 bound proteins are eluted with a 200-500 mM NaCl gradient. Fractions are analyzed by SDS-PAGE followed by immunoblotting with an Ab for hsp110, as described by Wang *et al.*, 1999, J. Immunol. 162:3378. Pooled fractions containing hsp110 are concentrated by Centriplus (Amicon, Beverly, MA) and applied to a Superose 12 column (Pharmacia). Proteins are eluted by 40 mM Tris-HCl, pH 8.0; 150 mM NaCl; and 15 mM 2-ME with a  
15 flow rate of 0.2 ml/min.

#### 4.2.6. Preparation and Purification of grp170-peptide Complexes

A procedure, described by Wang *et al.*, 2001, J. Immunol. 166(1):490-7, that  
20 can be used, presented by way of example and not limitation, is as follows:

A pellet (40-60 ml) of cell or tissue, *e.g.*, tumor cell tissue, is homogenized in 5 vol of hypotonic buffer (30 mM sodium bicarbonate, pH7.2, and protease inhibitors) by Dounce homogenization. The lysate is centrifuged at 4,500  $\times$  g and then 100,000  $\times$  g for 2 hours. If the cells or tissues are of hepatic origin, the resulting supernatant is was first  
25 applied to a blue Sepharose column (Pharmacia) to remove albumin. Otherwise, the resulting supernatant is applied to a Con A-Sepharose column (Pharmacia Biotech, Piscataway, NJ) previously equilibrated with binding buffer (20mM Tris-HCl, pH 7.5; 100mM NaCl; 1mM MgCl<sub>2</sub>; 1 mM CaCl<sub>2</sub>; 1 mM MnCl<sub>2</sub>; and 15 mM 2-ME). The bound proteins are eluted with binding buffer containing 15%  $\alpha$ -D-o-methylmannoside (Sigma, St. Louis, MO).  
30

Con A-Sepharose-bound material is first dialyzed against 20 mM Tris-HCl, pH 7.5, and 150 mM NaCl and then applied to a Mono Q column and eluted by a 150 to 400 mM NaCl gradient. Pooled fractions are concentrated and applied on the Superose 12 column (Pharmacia). Fractions containing homogeneous grp170 are collected.

35

#### 4.2.7. Infectious Disease

In an alternative embodiment wherein it is desired to treat a patient having an

infectious disease, the above-described methods are used to isolate  $\alpha$ 2M or hsp -antigenic molecule complexes from cells infected with an infectious organism, *e.g.*, of a cell line or from a patient. Such infectious organisms include but are not limited to, viruses, bacteria, protozoa, fungi, and parasites as described in detail in the Target Infectious Diseases

5 Section below.

#### 4.2.8. Autoimmune Disease

In an another alternative embodiment wherein it is desired to treat a patient having an autoimmune disease, the above-described methods are used to isolate  $\alpha$ 2M or hsp  
10 -antigenic molecule complexes from an autologous, allogeneic, or other cell line expressing an antigenic molecules of interest.

#### 4.2.9. Neurodegenerative and Amyloid Diseases

In an another alternative embodiment wherein it is desired to treat a patient  
15 having a neurodegenerative or amyloid disease, the above-described methods are used to isolate  $\alpha$ 2M or hsp-antigenic molecule complexes from an autologous, allogeneic, or other cell line expressing an antigenic molecules of interest. Examples of suitable cells and tissues from which such hsp-antigenic molecule complexes can be isolated are described in U.S. patent application no. 09/489,215, which is incorporated by reference herein in its  
20 entirety.

#### 4.2.10. Peptides From $\alpha$ 2M or hsp-Peptide Complexes

Antigenic molecules (*e.g.* peptides) can be eluted from hsp-antigenic molecule complexes either in the presence of ATP or low pH. Antigenic molecules can be  
25 eluted from  $\alpha$ 2M-antigenic molecule complexes in the presence of low pH. These experimental conditions may be used to isolate peptides or non-peptide antigenic components from cells which may contain potentially useful antigenic determinants. Once isolated, the amino acid sequence of an antigenic peptide may be determined using conventional amino acid sequencing methodologies. Antigenic molecules can then be  
30 produced by chemical synthesis or recombinant methods, purified, and complexed to hsps or  $\alpha$ 2M *in vitro*.

Thus, potentially immunogenic or antigenic peptides may be isolated from either endogenous stress protein-peptide complexes or endogenous MHC-peptide complexes for use subsequently as antigenic molecules, by complexing *in vitro* to hsps.

35 Briefly the complex of interest is centrifuged through a Centricon 10 assembly (Millipore) to remove any low molecular weight material loosely associated with

the complex. The large molecular weight fraction may be removed and analyzed by SDS-PAGE while the low molecular weight may be analyzed by HPLC as described below. In the ATP incubation protocol, the stress protein-peptide complex in the large molecular weight fraction is incubated with 10mM ATP for 30 minutes at room temperature. In the  
5 low pH protocol, acetic acid or trifluoroacetic acid (TFA) is added to the stress protein-peptide complex to give a final concentration of 10% (vol/vol) and the mixture incubated at room temperature or in a boiling water bath or any temperature in between, for 10 minutes (See, Udono *et al.*, 1993, J. Exp. Med. 178:1391-1396; Van Bleek *et al.*, 1990, Nature 348:213-216; and Li *et al.*, 1993, EMBO Journal 12:3143-3151).

10 The resulting samples are centrifuged through a Centricon 10 assembly as mentioned previously. The high and low molecular weight fractions are recovered. The remaining large molecular weight stress protein-peptide complexes can be reincubated with ATP or low pH to remove any remaining peptides.

The resulting lower molecular weight fractions are pooled, concentrated by  
15 evaporation and dissolved in 0.1% TFA. The dissolved material is then fractionated by reverse phase high pressure liquid chromatography (HPLC) using for example a VYDAC C18 reverse phase column equilibrated with 0.1% TFA. The bound material is then eluted at a flow rate of about 0.8 ml/min by developing the column with a linear gradient of 0 to 80% acetonitrile in 0.1% TFA. The elution of the peptides can be monitored by OD<sub>210</sub> and  
20 the fractions containing the peptides collected.

#### 4.2.11. Peptides From MHC-Peptide Complexes

The isolation of potentially immunogenic peptides from MHC molecules is well known in the art and so is not described in detail herein (See, Falk *et al.*, 1990, Nature  
25 348:248-251; Rotzsche *et al.*, 1990, Nature 348:252-254; Elliott *et al.*, 1990, Nature 348:191-197; Falk *et al.*, 1991, Nature 351:290-296; Demotz *et al.*, 1989, Nature 343:682-684; Rotzsche *et al.*, 1990, Science 249:283-287, the disclosures of which are incorporated herein by reference).

Briefly, MHC-peptide complexes may be isolated by a conventional  
30 immunoaffinity procedure. The peptides then may be eluted from the MHC-peptide complex by incubating the complexes in the presence of about 0.1% TFA in acetonitrile. The eluted peptides may be fractionated and purified by reverse phase HPLC, as before.

The amino acid sequences of the eluted peptides may be determined either by manual or automated amino acid sequencing techniques well known in the art. Once the  
35 amino acid sequence of a potentially protective peptide has been determined the peptide may be synthesized in any desired amount using conventional peptide synthesis or other



protocols well known in the art.

Peptides having the same amino acid sequence as those isolated above may be synthesized by solid-phase peptide synthesis using procedures similar to those described by Merrifield, 1963, J. Am. Chem. Soc. 85:2149. During synthesis, N- $\alpha$ -protected amino acids having protected side chains are added stepwise to a growing polypeptide chain linked by its C-terminal and to an insoluble polymeric support *i.e.*, polystyrene beads. The peptides are synthesized by linking an amino group of an N- $\alpha$ -deprotected amino acid to an  $\alpha$ -carboxy group of an N- $\alpha$ -protected amino acid that has been activated by reacting it with a reagent such as dicyclohexylcarbodiimide. The attachment of a free amino group to the activated carboxyl leads to peptide bond formation. The most commonly used N- $\alpha$ -protecting groups include Boc which is acid labile and Fmoc which is base labile.

Briefly, the C-terminal N- $\alpha$ -protected amino acid is first attached to the polystyrene beads. The N- $\alpha$ -protecting group is then removed. The deprotected  $\alpha$ -amino group is coupled to the activated  $\alpha$ -carboxylate group of the next N- $\alpha$ -protected amino acid. The process is repeated until the desired peptide is synthesized. The resulting peptides are then cleaved from the insoluble polymer support and the amino acid side chains deprotected. Longer peptides can be derived by condensation of protected peptide fragments. Details of appropriate chemistries, resins, protecting groups, protected amino acids and reagents are well known in the art and so are not discussed in detail herein (*See*, Atherton *et al.*, 1989, *Solid Phase Peptide Synthesis: A Practical Approach*, IRL Press, and Bodanszky, 1993, *Peptide Chemistry, A Practical Textbook*, 2nd Ed., Springer-Verlag).

Purification of the resulting peptides is accomplished using conventional procedures, such as preparative HPLC using gel permeation, partition and/or ion exchange chromatography. The choice of appropriate matrices and buffers are well known in the art and so are not described in detail herein.

#### 4.2.12. Exogenous Antigenic Molecules

Any antigen, or an antigenic portion thereof, can be selected for use as an antigenic molecule for complexing to hsp's, MHCs, or  $\alpha$ 2M. For example, antigens may be selected from among those known in the art or determined by immunoassay to be able to bind to antibody or MHC molecules (antigenicity) or generate an immune response (immunogenicity). To determine immunogenicity or antigenicity by detecting binding to antibody, various immunoassays known in the art can be used, including but not limited to competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitin reactions,

immunodiffusion assays, *in vivo* immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, immunoprecipitation reactions, agglutination assays (*e.g.*, gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, *etc.* In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labelled. Many means are known in the art for detecting binding in an immunoassay and are envisioned for use. In one embodiment for detecting immunogenicity, T cell-mediated responses can be assayed by standard methods, *e.g.*, *in vitro* cytotoxicity assays or *in vivo* delayed-type hypersensitivity assays.

Potentially useful antigens or derivatives thereof for use as antigenic molecules can also be identified by various criteria, such as the antigen's involvement in neutralization of a pathogen's infectivity (wherein it is desired to treat or prevent infection by such a pathogen) (Norrby, 1985, *Summary*, in Vaccines 85, Lerner *et al.* (eds.), Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, pp. 388-389), type or group specificity, recognition by patients' antisera or immune cells, and/or the demonstration of protective effects of antisera or immune cells specific for the antigen. In addition, where it is desired to treat or prevent a disease caused by pathogen, the antigen's encoded epitope should preferably display a small or no degree of antigenic variation in time or amongst different isolates of the same pathogen.

Preferably, where it is desired to treat or prevent cancer, known tumor-specific antigens or fragments or derivatives thereof are used. For example, such tumor specific or tumor-associated antigens include but are not limited to KS 1/4 pan-carcinoma antigen (Perez and Walker, 1990, J. Immunol. 142:3662-3667; Bumal, 1988, Hybridoma 7(4):407-415); ovarian carcinoma antigen (CA125) (Yu *et al.*, 1991, Cancer Res. 51(2):468-475); prostatic acid phosphate (Tailer *et al.*, 1990, Nucl. Acids Res. 18(16):4928); prostate specific antigen (Henttu and Vihko, 1989, Biochem. Biophys. Res. Comm. 160(2):903-910; Israeli *et al.*, 1993, Cancer Res. 53:227-230); melanoma-associated antigen p97 (Estin *et al.*, 1989, J. Natl. Cancer Inst. 81(6):445-446); melanoma antigen gp75 (Vijayasardahl *et al.*, 1990, J. Exp. Med. 171(4):1375-1380); high molecular weight melanoma antigen (Natali *et al.*, 1987, Cancer 59:55-63) and prostate specific membrane antigen.

In a specific embodiment, an antigen or fragment or derivative thereof specific to a certain tumor is selected for complexing to hsp and subsequent administration to a patient having that tumor.

Preferably, where it is desired to treat or prevent viral diseases, molecules

comprising epitopes of known viruses are used. For example, such antigenic epitopes may be prepared from viruses including, but not limited to, hepatitis type A, hepatitis type B, hepatitis type C, influenza, varicella, adenovirus, herpes simplex type I (HSV-I), herpes simplex type II (HSV-II), rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, papova virus, cytomegalovirus, echinovirus, arbovirus, huntavirus, coxsackie virus, mumps virus, measles virus, rubella virus, polio virus, human immunodeficiency virus type I (HIV-I), and human immunodeficiency virus type II (HIV-II).

Preferably, where it is desired to treat or prevent bacterial infections, molecules comprising epitopes of known bacteria are used. For example, such antigenic epitopes may be prepared from bacteria including, but not limited to, mycobacteria rickettsia, mycoplasma, neisseria and legionella.

Preferably, where it is desired to treat or prevent protozoal infections, molecules comprising epitopes of known protozoa are used. For example, such antigenic epitopes may be prepared from protozoa including, but not limited to, leishmania, kokzidioa, and trypanosoma.

Preferably, where it is desired to treat or prevent parasitic infections, molecules comprising epitopes of known parasites are used. For example, such antigenic epitopes may be from parasites including, but not limited to, chlamydia and rickettsia.

Preferably, where it is desired to treat or prevent neurodegenerative or amyloid diseases, molecules comprising epitopes of antigenic molecules associated with neurodegenerative diseases, or epitopes of antigenic molecules associated with amyloid diseases, including but not limited to fibril peptides or proteins, are used. For example, such neurodegenerative disease-associated antigenic molecules may be molecules associated with Alzheimer's Disease, age-related loss of cognitive function, senile dementia, Parkinson's disease, amyotrophic lateral sclerosis, Wilson's Disease, cerebral palsy, progressive supranuclear palsy, Guam disease, Lewy body dementia, prion diseases, spongiform encephalopathies, Creutzfeldt-Jakob disease, polyglutamine diseases, Huntington's disease, myotonic dystrophy, Freidrich's ataxia, ataxia, Gilles de la Tourette's syndrome, seizure disorders, epilepsy, chronic seizure disorder, stroke, brain trauma, spinal cord trauma, AIDS dementia, alcoholism, autism, retinal ischemia, glaucoma, autonomic function disorder, hypertension, neuropsychiatric disorder, schizophrenia, or schizoaffective disorder. Antigenic molecules that are suitable for *in vitro* complexing methods are disclosed in U.S. application no. 09/489,216, which is incorporated by reference herein in its entirety, and include, but are not limited to,  $\beta$ -amyloid, an oligomeric A $\beta$  complex, an ApoE4-A $\beta$  complex, tau protein, a mutant amyloid precursor, a mutant of presenillin,  $\alpha$ -

synuclein, a prion protein, or an antigenic fragment of any of the foregoing proteins.

Amyloid disease associated antigenic molecules may be molecules associated with diseases characterized by the extracellular deposition of protein and/or peptide fibrils which form amyloid deposits or plaques, including but not limited to type II diabetes and amyloidoses associated with chronic inflammatory or infectious disease states and malignant neoplasms, *e.g.*, myeloma. Certain amyloid disease such as Alzheimer's disease and prion diseases, *e.g.*, Creutzfeldt Jacob disease, are neurodegenerative diseases.

#### 4.2.13. *In Vitro* Production of hsp-Antigenic Molecule Complexes

In an embodiment in which complexes of hsps and the peptides with which they are endogenously associated *in vivo* are not employed, complexes of hsps to antigenic molecules are produced *in vitro*. As will be appreciated by those skilled in the art, the peptides either isolated by the aforementioned procedures or chemically synthesized or recombinantly produced may be reconstituted with a variety of purified natural or recombinant stress proteins *in vitro* to generate immunogenic non-covalent stress protein-antigenic molecule complexes. Alternatively, exogenous antigens or antigenic or immunogenic fragments or derivatives thereof can be complexed to stress proteins for use in the immunotherapeutic or prophylactic vaccines of the invention. A preferred, exemplary protocol for complexing a stress protein and an antigenic molecule *in vitro* is discussed below.

Prior to complexing, the hsps are pretreated with ATP or low pH to remove any peptides that may be associated with the hsp of interest. When the ATP procedure is used, excess ATP is removed from the preparation by the addition of apyranase as described by Levy *et al.*, 1991, *Cell* 67:265-274. When the low pH procedure is used, the buffer is readjusted to neutral pH by the addition of pH modifying reagents.

The antigenic molecules (1 $\mu$ g) and the pretreated hsp (9 $\mu$ g) are admixed to give an approximately 5 antigenic molecule: 1 stress protein molar ratio. Then, the mixture is incubated for 15 minutes to 3 hours at 4° to 45°C in a suitable binding buffer such as one containing 20mM sodium phosphate, pH 7.2, 350mM NaCl, 3mM MgCl<sub>2</sub> and 1mM phenyl methyl sulfonyl fluoride (PMSF). The preparations are centrifuged through a Centricon 10 assembly (Millipore) to remove any unbound peptide. The association of the peptides with the stress proteins can be assayed by SDS-PAGE. This is the preferred method for *in vitro* complexing of peptides isolated from MHC-peptide complexes of peptides disassociated from endogenous hsp-peptide complexes.

In an alternative embodiment of the invention, preferred for producing complexes of hsp70 to exogenous antigenic molecules such as proteins, 5-10 micrograms of

purified hsp is incubated with equimolar quantities of the antigenic molecule in 20mM sodium phosphate buffer pH 7.5, 0.5M NaCl, 3mM MgCl<sub>2</sub> and 1mM ADP in a volume of 100 microliter at 37°C for 1 hr. This incubation mixture is further diluted to 1ml in phosphate-buffered saline.

- 5 In an alternative embodiment of the invention, preferred for producing complexes of gp96 or hsp90 to peptides, 5-10 micrograms of purified gp96 or hsp90 is incubated with equimolar or excess quantities of the antigenic peptide in a suitable buffer such as one containing 20mM sodium phosphate buffer pH 7.5, 0.5M NaCl, 3mM MgCl<sub>2</sub> at 60-65°C for 5-20 min. This incubation mixture is allowed to cool to room temperature and  
10 centrifuged one or more times if necessary, through a Centricon 10 assembly (Millipore) to remove any unbound peptide.

- Following complexing, the immunogenic stress protein-antigenic molecule complexes can optionally be assayed *in vitro* using for example the mixed lymphocyte target cell assay (MLTC) described below. Once immunogenic complexes have been  
15 isolated they can be optionally characterized further in animal models using the preferred administration protocols and excipients discussed below.

#### 4.3. Sources of Saponins

- Any saponin or saponin preparation known in the art may be used in the  
20 compositions and methods of the invention. The term "saponin" as used herein includes glycosidic triterpenoid compounds which produce foam in aqueous solution and have hemolytic activity in most cases. The invention encompasses the use of saponins per se, as well as natural and pharmaceutically acceptable salts and pharmaceutically acceptable derivatives thereof. The term "saponin" also embodies biologically active fragments  
25 thereof. The term "saponin" also encompasses chemically modified saponins. In other embodiments of the invention, the term "saponin" covers mixtures of saponins. Preferably, the mixture of saponins comprises two or more substantially pure saponins. More preferably, the two or more substantially pure saponins are from *Quillaja saponaria* in doses that are otherwise suboptimal for the individual saponins. In a particularly preferred  
30 embodiment, the combination of saponins consists essentially of two or substantially pure saponins QS-7 and QS-21 or, in other particularly preferred embodiments, QS-7 and QS-21-V1 or QS-7 and QS-21-V2, as described in U.S. Patent No. 6,231, 859, which is herein incorporated by reference in its entirety. As used herein, "substantially pure" means  
substantially free from compounds normally associated with the saponin in its natural state  
35 and exhibiting constant and reproducible chromatographic response, elution profiles, and biologic activity. The term "substantially pure" is not meant to exclude artificial or

synthetic mixtures of the saponin with other compounds. A number of non-limiting examples of saponins and their methods of preparation are provided below.

Several embodiments of the invention relate to pharmaceutical compositions comprising either  $\alpha$ 2M or an hsp, a first antigen that will elicit a specific immune response to a cancer, infectious disease or neurodegenerative or amyloid disease, and a saponin adjuvant. The compositions of the invention comprising saponins can be formulated in a variety of ways to produce pharmaceutically effective compositions capable of eliciting an immune response in an individual to whom the composition is administered.

Additional embodiments of the invention relate to pharmaceutical compositions comprising either  $\alpha$ 2M or an hsp, optionally a peptide (which need not be antigenic), and a saponin adjuvant, for the prevention or treatment of an autoimmune disorder. These compositions are formulated to be capable of enhancing the suppression of an immune response in an individual to whom the composition is administered.

Saponins suitable for use in an individual are soluble in aqueous solution and can be reconstituted from lyophilized or dried saponins. Specific saponins useful for the present invention include, but are not limited to, the "Quil-A" adjuvant preparation sold by Superfos of Norway, and the chromatographic fractions with adjuvant activity that are described in U.S. Patent Nos. 5,057,540 and 5,583,112, particularly fractions QS-21 (also referred to in the patents as QA-21) and QS-7.

Also useful in the methods and compositions of the present invention are chemically modified saponins that retain adjuvant activity. According to Kensil *et al.*, U.S. Patent No. 5,583,112, the contents of which are fully incorporated by reference herein, the carboxyl group on the glucuronic acid of saponins from *Quillaja saponaria* Molina can be conjugated to a protein, a peptide, or a small molecule containing a primary amine. According to Higuchi *et al.*, 1987, *Phytochemistry* 26:229, saponins from *Quillaja saponaria* may be deacylated by alkaline-catalyzed hydrolysis. According to Marciani *et al.*, U.S. Patent No. 5,977,081, the contents of which are fully incorporated by reference herein, the carboxyl group on the glucuronic acid of nonacylated or deacylated saponins from *Quillaja saponaria* may be conjugated to a lipid, fatty acid, polyethylene glycol, or terpene.

Alternatively, an active fragment or synthetically modified derivative of a fragment or a native saponin(s) may be utilized, such as those described in Soltysik *et al.*, 1995, *Vaccine* 13(15):1403-1410; Marciani *et al.*, 2000, *Vaccine* 18:3141-3151. Such modifications include but are not limited to removals or substitutions of saccharide residues, addition of saccharide residues, and removal, substitution and/or addition of acyl chains.

The methods and compositions of the present invention may also employ

saponins isolated from plant species other than *Quillaja*, such as *Gypsophila* or *Saponaria officinalis*.

In certain embodiments of the invention, compositions of the invention comprises saponins in combination with excipients. Preferably, the saponin is QS-21 and  
5 the excipients are selected from nonionic surfactants, polyvinyl pyrrolidone, human serum albumin, and various unmodified and derivatized cyclodextrins. More preferably, in these embodiments, the nonionic surfactants are selected from Polysorbate 20, Polysorbate-40, Polysorbate-60, and Polysorbate-80. The polyvinyl pyrrolidone may preferably be Plasdone C15, a pharmaceutical grade of polyvinyl pyrrolidone. Preferred cyclodextrins are  
10 hydroxypropyl- $\beta$ -cyclodextrin, hydroxypropyl- $\gamma$ -cyclodextrin, and methyl- $\beta$ -cyclodextrin.

There are multiple acceptable techniques for extraction and isolation of saponins from *Quillaja saponaria* Molina bark. Acceptable procedures for purifying the saponins of the present invention from *Quillaja saponaria* Molina bark, measuring the saponins for immune adjuvant activity, and characterizing the substantially pure saponins  
15 are disclosed in U.S. Pat. Nos. 5,057,540 and 5,583,112.

Aqueous extracts of *Quillaja saponaria* bark are also available commercially. These are dark brown, foamy extracts that contain many compounds (tannins, polyphenolics, saponins) that can be analyzed by a method such as reversed phase HPLC.

20 An example of a reversed phase HPLC analysis of a typical bark extract that is suitable for purification of saponins is shown in FIG. 1 of U.S. Patent No. 6,231,859, which is incorporated herein in its entirety.

Partial purification to enrich the saponin fraction and to remove the majority of tannins and polyphenolics can be accomplished by dialysis of the extract against water  
25 through a 10,000 molecular weight membrane or ultrafiltration. The saponin fraction is retained.

Alternatively, an aqueous saponin extract can be pretreated with polyvinylpyrrolidone to remove high molecular weight tannins and polyphenolics through absorption of these compounds.

30 Residual tannins and polyphenolics can then be removed from the saponin fraction by diafiltration against water. The saponin fraction, which forms micelles, is retained by ultrafiltration membranes of 10,000 to 30,000 molecular weight cutoff pore size. This yields a partially purified extract that consists predominantly of diverse saponins.

Separation of saponins can be accomplished by chromatography in organic  
35 solvents or organic solvent/water mixtures. A separation of saponins on silica was described in U.S. Pat. No. 5,057,540. This yields saponins of intermediate purity (enriched

in an individual saponin, but less than substantially pure).

Alternatively, other solvent systems on silica gel or the use of reverse phase chromatography can be used to accomplish the initial separation of saponins. This initial purification step can then typically be followed by reversed phase chromatography or  
5 similar HPLC step to purify the saponins to near homogeneity.

For example, saponin extract may be recovered from plant cell material freshly extracted from *Quillaja* trees. Dialyzed extract is then purified on an ion exchange column, e.g., the DE-52 type, followed by Sephadex G50 gel filtration. Ultrafiltration may be used instead of gel filtration. The purified saponin composition is then subjected to  
10 RP-HPLC analysis on a VYDAC C4 column, eluted with 30-45% acetonitrile in a 0.15% aqueous TFA-solution.

The substantially pure saponins useful in the present invention may also be isolated from fresh plant material consisting of substantially living cells as disclosed in WO 95/09179, or the previously described procedures.

15 The same procedure may be performed on plant cell material obtained by means of tissue culture or suspension cell culture. See, e.g., U.S. Patent No. 5,716,848, which is incorporated herein by reference in its entirety.

General guidance on the use of saponins, Quil-A, and QS-21 can be found in the referenced patents. The amount of saponin present in a pharmaceutically effective  
20 composition should contain about 0.1 to 5,000 micrograms or more of a saponin. The amount of saponin present in a pharmaceutically effective composition is more preferably from about 1 to about 1000 micrograms, more preferably from about 5 to 500 micrograms, and most preferably from about 10 to 100 micrograms. In certain specific embodiments, the amount of saponin present in a pharmaceutical composition of the invention is 1, 2, 3, 5, 10,  
25 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 125 or 150 micrograms.

In certain embodiments of the present invention, a composition of the invention comprises a saponin covalently linked to an antigenic molecule. Covalent linkage of antigens to saponins is fully described in U.S. Patent No. 5,583,112 at column 9, lines 4-67 and column 10, line 1-37. Additionally, one of skill in the art will be able to establish  
30 crosslinking conditions without undue experimentation such that covalent linkage of saponin to antigen and hsp to antigen may be carried out simultaneously.

Formulations of pharmaceutical compositions comprising saponins and procedures for their manufacture can be found in the literature and in the U.S. Patents incorporated by reference into this Description. Saponin formulations referred to are  
35 provided herein as nonlimiting examples.

In U.S. Patent No. 5,583,112 at column 22, lines 11-17, a pharmaceutically



effective composition for intradermal administration was made by reconstituting lyophilized "Quil A", a crude saponin mixture, into a phosphate buffered saline (PBS) solution and mixed with a solution containing 10 micrograms of Bovine Serum Albumin (BSA) to achieve a final volume of 200 microliters for intradermal injection. The effective amount of  
5 "Quil A" was found to be approximately 30-77 micrograms of "Quil A" by dry weight.

In U.S. Patent No. 5,583,112 at column 23, lines 35-38, a pharmaceutically effective composition for subcutaneous administration was made by mixing a stock saline solution of ovalbumin and 10 micrograms of QS-21.

Also in U.S. Patent No. 5,583,112 at Column 23, lines 35-38, a  
10 pharmaceutically effective composition for subcutaneous administration was made by chemically crosslinking QS-21 to lysozyme as described in Example 18 of the patent and resuspending lyophilized QS-21/lysozyme conjugate into 200 microliters of PBS (pH 7) for a final concentration of 10 micrograms of lysozyme and 1.6 micrograms of QS-21.

In a paper by Wu *et al.* (1994, Cellular Immunology 154:393-406), a  
15 pharmaceutically effective composition for subcutaneous or intraperitoneal administration is disclosed containing 25 micrograms ovalbumin absorbed to 250 micrograms of  $Al(OH)_3$  and 20 micrograms of QS-21 per immunization dose.

In another paper by Wu *et al.* (1994, J. Immun. 148(5):1519-1525), a pharmaceutically effective composition for immunization is disclosed containing 25  
20 micrograms of a truncated recombinant HIV-1 envelope protein absorbed to 250 micrograms of  $Al(OH)_3$  and 10 micrograms of QS-21 in a sterile saline per immunization dose.

As a last non-limiting example, a pharmaceutically effective vaccine has recently been tested in human patients containing 5-500 micrograms of a synthetic  
25 nonapeptide and 100 micrograms of QS-21 in 500 microliters PBS (pH 7.4) per intradermally-administered dose (Lewis *et al.*, 2000, Int. J. Cancer 87(3):391-398).

The optimum amount of a specific saponin for use with a specific composition of the invention may vary. Optimization of the specific saponin amount for a given composition is, as demonstrated by the examples cited above, well within the purview  
30 of the skilled artisan.

#### 4.4. Determination of Immunogenicity

The compositions of the invention can be assayed for immunogenicity using any method known in the art. In certain embodiments, the antigenicity of the individual  
35 components of a composition of the invention can be assayed alone or in combination, and compared to the antigenicity of the composition as whole, to determine the enhancement of

the immunogenicity of the composition over the immunogenicity of its individual components. For example, the immunogenicity of an hsp-antigenic molecule complex can be assayed and compared to immunogenicity of a composition comprising the hsp-antigenic molecule and a saponin. Preferably, a composition of the invention is more immunogenic  
5 than the immunogenicity of its individual components, and, more preferably is synergistically more immunogenic than the immunogenicity of its individual components. In one specific embodiment, an immunogenic composition of the invention comprises sub-immunogenic amounts of its individual components, *i.e.*, comprises, for example, an hsp-antigenic molecule in an amount that is in itself non-immunogenic, but is immunogenic  
10 when combined with a saponin according to the disclosed methods.

By way of example but not limitation, any one of the following procedures can be used. In a preferred embodiment, the ELISPOT assay is used (see, *infra*, Section 4.4.4).

#### 15 4.4.1. The MLTC Assay

Briefly, mice are injected with an amount of an  $\alpha 2M$  or hsp-antigenic molecule complex, using any convenient route of administration. As a negative control, other mice are injected with  $\alpha 2M$  or hsp-control complexes (*e.g.*  $\alpha 2M$  or hsp stripped of antigenic molecules or complexed with non-antigenic molecules). Cells known to contain  
20 specific antigens, *e.g.* tumor cells or cells infected with an agent of an infectious disease, may act as a positive control for the assay. The mice are injected twice, 7-10 days apart. Ten days after the last immunization, the spleens are removed and the lymphocytes released. The released lymphocytes may be re-stimulated subsequently *in vitro* by the addition of dead cells that expressed the antigen of interest.

25 For example,  $8 \times 10^6$  immune spleen cells may be stimulated with  $4 \times 10^4$  mitomycin C treated or  $\gamma$ -irradiated (5-10,000 rads) cells containing the antigen of interest (or cells transfected with an appropriate gene, as the case may be) in 3ml RPMI medium containing 10% fetal calf serum. In certain cases 33% secondary mixed lymphocyte culture supernatant may be included in the culture medium as a source of T cell growth factors  
30 (*See*, Glasebrook *et al.*, 1980, J. Exp. Med. 151:876). To test the primary cytotoxic T cell response after immunization, spleen cells may be cultured without stimulation. In some experiments spleen cells of the immunized mice may also be re-stimulated with antigenically distinct cells, to determine the specificity of the cytotoxic T cell response.

Six days later the cultures are tested for cytotoxicity in a 4 hour  $^{51}\text{Cr}$ -release  
35 assay (*See*, Palladino *et al.*, 1987, Cancer Res. 47:5074-5079 and Blachere *et al.*, 1993, J. Immunotherapy 14:352-356). In this assay, the mixed lymphocyte culture is added to a

target cell suspension to give different effector:target (E:T) ratios (usually 1:1 to 40:1). The target cells are prelabelled by incubating  $1 \times 10^6$  target cells in culture medium containing 20 mCi  $^{51}\text{Cr}$ /ml for one hour at  $37^\circ\text{C}$ . The cells are washed three times following labeling. Each assay point (E:T ratio) is performed in triplicate and the appropriate controls

5 incorporated to measure spontaneous  $^{51}\text{Cr}$  release (no lymphocytes added to assay) and 100% release (cells lysed with detergent). After incubating the cell mixtures for 4 hours, the cells are pelleted by centrifugation at 200g for 5 minutes. The amount of  $^{51}\text{Cr}$  released into the supernatant is measured by a gamma counter. The percent cytotoxicity is measured as cpm in the test sample minus spontaneously released cpm divided by the total detergent

10 released cpm minus spontaneously released cpm.

In order to block the MHC class I cascade a concentrated hybridoma supernatant derived from K-44 hybridoma cells (an anti-MHC class I hybridoma) is added to the test samples to a final concentration of 12.5%.

#### 15 4.4.2. CD4+ T Cell Proliferation Assay

Primary T cells are obtained from spleen, fresh blood, or CSF and purified by centrifugation using FICOLL-PAQUE PLUS (Pharmacia, Uppsala, Sweden) essentially as described by Kruse and Sebal, 1992, EMBO J. 11: 3237-3244. The peripheral blood mononuclear cells are incubated for 7-10 days with a lysate of cells expressing an antigenic

20 molecule. Antigen presenting cells may, optionally be added to the culture 24 to 48 hours prior to the assay, in order to process and present the antigen in the lysate. The cells are then harvested by centrifugation, and washed in RPMI 1640 media (GibcoBRL, Gaithersburg, Md.).  $5 \times 10^4$  activated T cells/well are in RPMI 1640 media containing 10% fetal bovine serum, 10 mM HEPES, pH 7.5, 2 mM L-glutamine, 100 units/ml penicillin G,

25 and 100  $\mu\text{g}/\text{ml}$  streptomycin sulphate in 96 well plates for 72 hrs at  $37^\circ\text{C}$ ., pulsed with 1  $\mu\text{Ci}$   $^3\text{H}$ -thymidine (DuPont NEN, Boston, Mass.)/well for 6 hrs, harvested, and radioactivity measured in a TOPCOUNT scintillation counter (Packard Instrument Co., Meriden,CT).

#### 4.4.3. Antibody Response Assay

30 In a certain embodiment of the invention, the immunogenicity of an  $\alpha 2\text{M}$  or hsp -antigenic molecule complex is determined by measuring antibodies produced in response to the vaccination with the complex. In one mode of the embodiment, microtitre plates (96-well Immuno Plate II, Nunc) are coated with 50  $\mu\text{l}/\text{well}$  of a 0.75  $\mu\text{g}/\text{ml}$  solution of a purified, non- $\alpha 2\text{M}$  or hsp-complexed form of the antigenic molecule used in the

35 vaccine (e.g. A $\beta$ 42) in PBS at  $4^\circ\text{C}$  for 16 hours and at  $20^\circ\text{C}$  for 1 hour. The wells are emptied and blocked with 200  $\mu\text{l}$  PBS-T-BSA (PBS containing 0.05% (v/v) TWEEN 20

and 1% (w/v) bovine serum albumin) per well at 20°C for 1 hour, then washed 3 times with PBS-T. Fifty µl/well of plasma or CSF from a vaccinated animal (such as a model mouse or a human patient) is applied at 20°C for 1 hour, and the plates are washed 3 times with PBS-T. The anti-peptide antibody activity is then measured calorimetrically after incubating  
5 at 20°C for 1 hour with 50µl/well of sheep anti-mouse or anti-human immunoglobulin, as appropriate, conjugated with horseradish peroxidase (Amersham) diluted 1:1,500 in PBS-T-BSA and (after 3 further PBS-T washes as above) with 50 µl of an o-phenylene diamine (OPD)-H<sub>2</sub>O<sub>2</sub> substrate solution. The reaction is stopped with 150 µl of 2M H<sub>2</sub>SO<sub>4</sub> after 5 minutes and absorbance is determined in a Kontron SLT-210 photometer (SLT  
10 Lab-instr., Zurich, Switzerland) at 492 nm (ref. 620 nm).

#### 4.4.4. Cytokine Detection Assays

The CD4+ T cell proliferative response to the compositions of the invention may be measured by detection and quantitation of the levels of specific cytokines. In one  
15 embodiment, for example, intracellular cytokines may be measured using an IFN-γ detection assay to test for immunogenicity of a complex of the invention. In an example of this method, peripheral blood mononuclear cells from a subject treated with a composition of the invention are stimulated with peptide antigens of a given tumor or with peptide  
antigens of an agent of infectious disease. Cells are then stained with T cell-specific labeled  
20 antibodies detectable by flow cytometry, for example FITC-conjugated anti-CD8 and PerCP-labeled anti-CD4 antibodies. After washing, cells are fixed, permeabilized, and reacted with dye-labeled antibodies reactive with human IFN-γ (PE- anti-IFN-γ). Samples are analyzed by flow cytometry using standard techniques.

Alternatively, a filter immunoassay, the enzyme-linked immunospot assay  
25 (ELISPOT) assay, may be used to detect specific cytokines surrounding a T cell. In one embodiment, for example, a nitrocellulose-backed microtiter plate is coated with a purified cytokine-specific primary antibody, *i.e.*, anti-IFN-γ, and the plate is blocked to avoid background due to nonspecific binding of other proteins. A sample of mononuclear blood cells containing cytokine-secreting cells is obtained from a subject treated with a  
30 composition of the invention, which sample is diluted onto the wells of the microtitre plate. A label, *e.g.*, biotin-labeled, secondary anti-cytokine antibody, is added. The antibody-cytokine complex can then be detected, *i.e.* by enzyme-conjugated streptavidin – cytokine-secreting cells will appear as “spots” by visual, microscopic, or electronic detection methods.

35

#### 4.4.5. Tetramer Assay

In another embodiment, the "tetramer staining" assay (Altman et al., 1996, Science 274: 94-96) may be used to identify antigen-specific T-cells. For example, in one embodiment, an MHC molecule containing a specific peptide antigen, such as a tumor-specific antigen, is multimerized to make soluble peptide tetramers and labeled, for example, by complexing to streptavidin. The MHC-peptide antigen complex is then mixed with a population of T cells obtained from a subject treated with a composition of the invention. Biotin is then used to stain T cells which express the antigen of interest, *i.e.*, the tumor-specific antigen.

#### 4.5. Recombinant Expression of hsps or $\alpha$ 2M

In certain embodiments of the invention, the compositions and methods comprise recombinant hsps, alone or complexed to antigenic molecules, or hsp-antigenic molecule complexes prepared from cells that express enhanced levels of hsps through recombinant means. In other embodiments of the invention, the compositions and methods comprise recombinant  $\alpha$ 2M or  $\alpha$ 2M-antigenic molecule complexes comprising recombinant  $\alpha$ 2M. In this regard, any method known to the skilled artisan may be used for obtaining and manipulating recombinant hsp or  $\alpha$ 2M sequences. Described below are non-limiting examples of such methods.

##### 4.5.1. hsp Sequences

Amino acid sequences and nucleotide sequences of many hsps are generally available in sequence databases, such as GenBank. Computer programs, such as Entrez, can be used to browse the database, and retrieve any amino acid sequence and genetic sequence data of interest by accession number. These databases can also be searched to identify sequences with various degrees of similarities to a query sequence using programs, such as FASTA and BLAST, which rank the similar sequences by alignment scores and statistics. Such nucleotide sequences of non-limiting examples of hsps that can be used for the compositions, methods, and for preparation of the hsp-antigenic molecule complexes of the invention are as follows: human hsp70, Genbank Accession No.M24743, Hunt *et al.*, 1995, Proc. Natl. Acad. Sci. U.S.A., 82: 6455-6489; human hsp90, Genbank Accession No.X15183, Yamazaki *et al.*, Nucl. Acids Res. 17: 7108; human gp96, Genbank Accession No.X15187, Maki *et al.*, 1990, Proc. Natl. Acad. Sci. U.S.A. 87: 5658-5562; human BiP, Genbank Accession No.M19645, Ting *et al.*, 1988, DNA 7: 275-286; human hsp27, Genbank Accession No.M24743, Hickey *et al.*, 1986, Nucleic Acids Res. 14: 4127-45; mouse hsp70, Genbank Accession No.M35021, Hunt *et al.*, 1990, Gene 87: 199-204; mouse

gp96, Genbank Accession No.M16370, Srivastava *et al.*, 1987, Proc. Natl. Acad. Sci. U.S.A. 85: 3807-3811; and mouse BiP, Genbank Accession No.U16277, Haas *et al.*, 1988, Proc. Natl. Acad. Sci. U.S.A. 85: 2250-2254. Due to the degeneracy of the genetic code, the term "hsp gene", as used herein, refers not only to the naturally occurring nucleotide  
5 sequence but also encompasses all the other degenerate DNA sequences that encode the hsp.

Once the nucleotide sequence encoding the hsp of choice has been identified, the nucleotide sequence, or a fragment thereof, can be obtained (*e.g.* from commercial sources or by PCR as described below) and cloned into an expression vector for  
10 recombinant expression. The expression vector can then be introduced into a host cell for propagation of the hsp. Methods for recombinant production of hsps are quite well known, as exemplified herein.

The DNA may be obtained by DNA amplification or molecular cloning directly from a tissue, cell culture, or cloned DNA (*e.g.*, a DNA "library") using standard  
15 molecular biology techniques (see *e.g.*, Methods in Enzymology, 1987, volume 154, Academic Press; Sambrook *et al.* 1989, Molecular Cloning - A Laboratory Manual, 2nd Edition, Cold Spring Harbor Press, New York; and Current Protocols in Molecular Biology, Ausubel *et al.* (eds.), Greene Publishing Associates and Wiley Interscience, New York, each of which is incorporated herein by reference in its entirety). Clones derived from genomic  
20 DNA may contain regulatory and intron DNA regions in addition to coding regions; clones derived from cDNA will contain only exon sequences. Whatever the source, the hsp gene should be cloned into a suitable vector for propagation of the gene.

In a preferred embodiment, DNA can be amplified from genomic or cDNA by polymerase chain reaction (PCR) amplification using primers designed from the known  
25 sequence of a related or homologous hsp. PCR is used to amplify the desired sequence in DNA clone or a genomic or cDNA library, prior to selection. PCR can be carried out, *e.g.*, by use of a thermal cycler and Taq polymerase (Gene Amp®). The polymerase chain reaction (PCR) is commonly used for obtaining genes or gene fragments of interest. For example, a nucleotide sequence encoding an hsp of any desired length can be generated  
30 using PCR primers that flank the nucleotide sequence encoding open reading frame. Alternatively, an hsp gene sequence can be cleaved at appropriate sites with restriction endonuclease(s) if such sites are available, releasing a fragment of DNA encoding the hsp gene. If convenient restriction sites are not available, they may be created in the appropriate positions by site-directed mutagenesis and/or DNA amplification methods known in the art  
35 (see, for example, Shankarappa *et al.*, 1992, PCR Method Appl. 1: 277-278). The DNA



fragment that encodes the hsp is then isolated, and ligated into an vector, care being taken to ensure that the proper translation readi

In an alternative embodiment, for the molecular cl  
genomic DNA, DNA fragments are generated to form a genomic  
5 sequences encoding related hsps are available and can be purified  
DNA fragments in the genomic DNA library may be screened by  
to a labeled probe (Benton and Davis, 1977, Science 196: 180; G  
1975, Proc. Natl. Acad. Sci. U.S.A. 72: 3961). Those DNA fragr  
homology to the probe will hybridize. It is also possible to identi  
10 by restriction enzyme digestion(s) and comparison of fragment si  
according to a known restriction map.

Alternatives to isolating the hsp genomic DNA inc  
to, chemically synthesizing the gene sequence itself from a know  
a cDNA to the mRNA which encodes the hsp. For example, RN<sub>2</sub>  
15 hsp gene can be isolated from cells which express the hsp. A cDI  
generated by methods known in the art and screened by methods,  
for screening a genomic DNA library. If an antibody to the hsp is  
identified by binding of a labeled antibody to the hsp-synthesizin

Other specific embodiments for the cloning of a n  
20 encoding an hsp, are presented as examples but not by way of lim  
specific embodiment, nucleotide sequences encoding an hsp can t  
by hybridization with a probe comprising a nucleotide sequence e  
conditions of low to medium stringency. By way of example and  
using such conditions of low stringency are as follows (see also S  
25 Proc. Natl. Acad. Sci. U.S.A. 78: 6789-6792). Filters containing I  
at 40°C in a solution containing 35% formamide, 5X SSC, 50 mM  
EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatur  
Hybridizations are carried out in the same solution with the follo  
0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm I  
30 dextran sulfate, and 5-20x10<sup>6</sup> cpm <sup>32</sup>P-labeled probe is used. Filt  
hybridization mixture for 18-20h at 40°C, and then washed for 1..  
containing 2X SSC, 25 mM Tris-HCl (pH7.4), 5mM EDTA, and  
solution is replaced with fresh solution and incubated an addition  
blotted dry and exposed for autoradiography. If necessary, filters  
35 time at 65-68°C and reexposed to film. Other conditions of low s  
used are well known in the art (e.g., as employed for cross-species



Any technique for mutagenesis known in the art can be used to alter individual nucleotides in a DNA sequence, for purpose of making changes in the expressed peptide sequence, or for creating/deleting restriction sites for further manipulations. Such techniques include but are not limited to: random mutagenesis, *in vitro* site-directed mutagenesis (Hutchinson *et al.*, 1985, 253: 6551), oligonucleotide-directed mutagenesis (Smith, 1985, A 463; Hill *et al.*, 1987, Methods Enzymol. 155: 558-568), PCR-based mutagenesis (Saiki *et al.*, 1989, Gene 77: 51-59), PCR-based megaprimer mutagenesis (Biotéchniques 8: 404-407), *etc.* Modifications can be confirmed, for example, by dideoxynucleotide DNA sequencing.

In certain embodiments, a nucleic acid encoding a polypeptide of choice is used to prepare the compositions and/or practice the method of the invention. Such a nucleic acid can be constructed by, *e.g.*, deleting an ER retention signal, KDEL. Optionally, the KDEL coding sequence can be replaced by a molecular tag, such as the Fc portion of murine IgG1, to facilitate purification of the hsp. U.S. Application No. 09/253,439, incorporated by reference, demonstrates that deletion of the ER retention signal of gp96 results in gp96-Ig peptide-complexes from transfected tumor cells, and that the deleted gp96 with murine IgG1 facilitated its detection by ELISA and its purification by affinity chromatography with the aid of Protein A.

#### 4.5.2. $\alpha$ 2M Sequences

$\alpha$ 2M polypeptides may be produced by recombinant DNA synthetic methods, or by enzymatic or chemical cleavage of native polypeptides. Described below are methods for producing such  $\alpha$ 2M polypeptides.

In various aspects, the invention relates to compositions and methods for producing  $\alpha$ 2M polypeptides. In various aspects, acid sequences of  $\alpha$ 2M, and fragments, derivatives, analogs, and variants of  $\alpha$ 2M are provided, as well as nucleic acids complementary to  $\alpha$ 2M hybridizing to such nucleic acids.

Any eukaryotic cell may serve as the nucleic acid source for the coding region of an  $\alpha$ 2M gene. Nucleic acid sequences encoding  $\alpha$ 2M are available from vertebrate, mammalian, as well as primate sources, including human.

Amino acid sequences and nucleotide sequences of  $\alpha$ 2M polypeptides are generally available in sequence databases, such as Genbank. Examples of  $\alpha$ 2M sequences that can be used for preparation of the compositions of the invention are as follows: Genbank Accession Nos. M11313, P01010, and P01011.

Kan *et al.*, 1985, Proc. Nat. Acad. Sci. U.S.A. 82: 2282-2286. In the genetic code, the term " $\alpha$ 2M gene", as used herein, refers not only to the occurring nucleotide sequence but also encompasses all the other sequences that encode an  $\alpha$ 2M polypeptide. Computer programs used to browse the database, and retrieve any amino acid sequence data of interest by accession number. These databases can also be used to search for sequences with various degrees of similarities to a query sequence. FASTA and BLAST, which rank the similar sequences by alignment. Nucleotide BLAST searches can be performed with the NBLAST program with wordlength = 12 to obtain nucleotide sequences homologous to the invention. BLAST protein searches can be performed with the BLAST program with wordlength = 3 to obtain amino acid sequences homologous to the invention. To obtain gapped alignments for comparison, Gapped BLAST can be utilized as described in Altschul *et al.*, 1997, Nuc. Res. 25:3389-3402. Alternatively, PSI-Blast can be used to perform searches which detects distant relationships between molecules (Altschul *et al.*, 1990). Utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the respective programs (*e.g.*, XBLAST and NBLAST) can be used (<http://www.ncbi.nlm.nih.gov>).

The DNA may be obtained by standard procedure, such as PCR amplification or molecular cloning directly from a tissue, cell culture, or a DNA "library". Clones derived from genomic DNA may contain non-coding DNA regions in addition to coding regions; clones derived from cDNA libraries contain only exon sequences. Whatever the source, the  $\alpha$ 2M gene should be confirmed by sequencing. The DNA may be used for propagation of the gene.

In a preferred embodiment, DNA can be amplified by polymerase chain reaction (PCR) amplification using primers flanking the sequence of a related or homologous  $\alpha$ 2M. PCR is used to amplify a DNA clone or a genomic or cDNA library, prior to selection. PCR can be performed by use of a thermal cycler and Taq polymerase (sold under the trade name of AmpliTaq, Perkin-Elmer). The DNA being amplified can include cDNA or genomic DNA flanked by oligonucleotide primers representing known nucleic acid sequences. Oligonucleotide primers used as primers in PCR. In a preferred aspect, the oligonucleotide primers are part of the  $\alpha$ 2M gene that is highly conserved between  $\alpha$ 2M genes. One can choose to synthesize several different degenerate primers, for example, to amplify the  $\alpha$ 2M gene. It is also possible to vary the stringency of hybridization conditions.

PCR reactions, to allow for greater or lesser degrees of nucleotide  
 between the known  $\alpha 2M$  nucleotide sequence and the nucleic acid  
 For cross species hybridization, low stringency conditions are preferred  
 hybridization, moderately stringent conditions are preferred. After  
 5 the sequence encoding an  $\alpha 2M$  may be cloned and sequenced. If the  
 region of the  $\alpha 2M$  gene being amplified is too large to be amplified  
 PCR covering the entire gene, preferably with overlapping regions,  
 the products of the PCR ligated together to form the entire coding sequence  
 if a segment of an  $\alpha 2M$  gene is amplified, that segment may be cloned  
 10 probe to isolate a complete cDNA or genomic clone.

In another embodiment, for the molecular cloning of  
 genomic DNA, DNA fragments are generated to form a genomic library  
 sequences encoding related  $\alpha 2Ms$  are available and can be purified  
 DNA fragments in the genomic DNA library may be screened by nucleic  
 15 to the labeled probe (Benton and Davis, 1977, Science 196:180; Gr  
 1975, Proc. Natl. Acad. Sci. U.S.A. 72:3961). Those DNA fragments with  
 homology to the probe will hybridize. It is also possible to identify  
 fragment by restriction enzyme digestion(s) and comparison of fragments  
 expected according to a known restriction map if such is available.

20 Alternatives to isolating the  $\alpha 2M$  genomic DNA include  
 to, chemically synthesizing the gene sequence itself from a known  
 cDNA to the mRNA which encodes  $\alpha 2M$ . For example, RNA for the  
 $\alpha 2M$  gene can be isolated from cells which express  $\alpha 2M$ . A cDNA  
 generated by methods known in the art and screened by methods, suitable  
 25 for screening a genomic DNA library. If an antibody to  $\alpha 2M$  is available  
 identified by binding of labeled antibody to the putatively  $\alpha 2M$  syn

Other specific embodiments for the cloning of a nucleic acid  
 encoding an  $\alpha 2M$ , are presented as examples but not by way of limitation.

In a specific embodiment, nucleotide sequences encoding  
 30 within a family can be identified and obtained by hybridization with  
 nucleotide sequence encoding  $\alpha 2M$  under conditions of low to medium

By way of example and not limitation, procedures useful for  
 low stringency are as follows (see also Shilo and Weinberg, 1981, J  
 USA 78:6789-6792). Filters containing DNA are pretreated for 6 hours  
 35 containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM  
 0.1% Ficoll, 1% BSA, and 500  $\mu$ g/ml denatured salmon sperm DNA

carried out in the same solution with the following modifications:  
 Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, 10% (wt/vol) (5-20 X 10<sup>6</sup> cpm <sup>32</sup>P-labeled probe is used. Filters are incubated in  
 18-20 h at 40°C, and then washed for 1.5 h at 55°C in a solution (5  
 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS. The wash  
 fresh solution and incubated an additional 1.5 h at 60°C. Filters are  
 exposed for autoradiography. If necessary, filters are washed for  
 and reexposed to film. Other conditions of low stringency which  
 known in the art (e.g., as employed for cross-species hybridization

10 An α2M gene fragment can be inserted into an app  
 and introduced into host cells so that many copies of the gene seq  
 large number of vector-host systems known in the art may be used  
 to, bacteriophages such as lambda derivatives, or plasmids such as  
 derivatives or the Bluescript vector (Stratagene).

15 Any technique for mutagenesis known in the art ca  
 individual nucleotides in a DNA sequence, for purpose of making  
 in the expressed peptide sequence, or for creating/deleting restricti  
 further manipulations. Such techniques include but are not limite  
 mutagenesis, *in vitro* site-directed mutagenesis (Hutchinson *et al.*,  
 20 253:6551), oligonucleotide-directed mutagenesis (Smith, 1985, A  
 463; Hill *et al.*, 1987, Methods Enzymol. 155:558-568), PCR-bas  
*et al.*, 1989, Gene 77:51-59), PCR-based megaprimer mutagenesis  
 Biotechniques, 8:404-407), *etc.* Modifications can be confirmed b  
 dideoxy DNA sequencing.

25 The polymerase chain reaction (PCR) is commonly  
 or gene fragments of interest. For example, a nucleotide sequence  
 polypeptide of any desired length can be generated using PCR pri  
 nucleotide sequence encoding α2M, or the peptide-binding domain  
 an α2M gene sequence can be cleaved at appropriate sites with res  
 30 if such sites are available, releasing a fragment of DNA encoding  
 binding domain thereof. If convenient restriction sites are not ava  
 created in the appropriate positions by site-directed mutagenesis a  
 methods known in the art (see, for example, Shankarappa *et al.*, 19  
 1:277-278). The DNA fragment that encodes α2M, or the peptide  
 35 is then isolated, and ligated into an appropriate expression vector,  
 that the proper translation reading frame is maintained.

Alpha (2) macroglobulin polypeptides may be expected to facilitate recovery and purification from the cells in which they are produced. For example, an  $\alpha 2M$  polypeptide may contain a signal sequence located at the N-terminal for translocation across the ER membrane for secretion into culture medium. Another  $\alpha 2M$  polypeptide may contain an affinity label, such as a fluorescent label, at the C-terminal. The  $\alpha 2M$  polypeptide not involved in binding antigenic peptide, is a separate polypeptide. The carboxyl terminal. The affinity label can be used to facilitate purification by binding to an affinity partner molecule.

Various methods for production of such fusion proteins are known in the art. The manipulations which result in their production can occur at the gene level, preferably at the gene level. For example, the cloned coding sequence for an  $\alpha 2M$  polypeptide may be modified by any of numerous recombinant DNA techniques (Sambrook *et al.*, 1990, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York; Ausubel *et al.*, 1994, Protocols in Molecular Biology, Greene Publishing Associates and Wiley Interscience, New York). It will be apparent from the following discussion that substitutions, deletions, or any combination thereof are introduced or combined with the nucleotide sequence encoding an  $\alpha 2M$  polypeptide.

In various embodiments, fusion proteins comprising an  $\alpha 2M$  polypeptide may be made using recombinant DNA techniques. For example, a fusion protein encoding an  $\alpha 2M$  polypeptide may be constructed by introducing the proper reading frame into a vector containing the sequence of the  $\alpha 2M$  polypeptide is expressed as a peptide-tagged fusion protein. The fusion protein may be recognized by specific binding partners, may be used for purification, and may be used for other purposes.

In a preferred embodiment, the affinity label is fused to the carboxyl terminal of  $\alpha 2M$ . The precise site at which the fusion is made is not critical. The optimal site can be determined by routine experimentation.

A variety of affinity labels known in the art may be used. For example, the affinity label may be limited to, the immunoglobulin constant regions, polyhistidine sequence, metal-chelate affinity chromatography, in Current Protocols in Molecular Biology, Ed. Ausubel *et al.*, Greene Publish. Assoc. & Wiley Interscience), (GST; Smith, 1993, Methods Mol. Cell Bio. 4:220-229), the *E. coli* (Guan *et al.*, 1987, Gene 67:21-30), and various cellulose binding proteins (Nos. 5,496,934; 5,202,247; 5,137,819; Tomme *et al.*, 1994, Protein Engineering). Other affinity labels may impart fluorescent properties to an  $\alpha 2M$  polypeptide.

of green fluorescent protein and the like. Other possible affinity sequences to which monoclonal antibodies are available, such as following well known examples, the FLAG epitope, the myc epitope, the influenza virus hemagglutinin (HA) epitope. Other affinity labels can be provided by specific binding partners and thus facilitate isolation by affinity chromatography using a partner which can be immobilized onto a solid support. Some affinity labels are derived from the novel structural properties, such as the ability to dimerize, of an  $\alpha 2M$  polypeptide. Dimerization of an  $\alpha 2M$  polypeptide with a bound peptide may interfere with the interaction between the  $\alpha 2M$  polypeptide and its partner in the complex presentation. These affinity labels are usually derived from protein homopolymers. Affinity labels such as the extracellular domains of CD28 (Lee *et al.*, 1988, J. Exp. Med. 168:1993-2005), or CD28 (Lee *et al.*, 1990, J. Exp. Med. 171:1003-1010), or portions of the immunoglobulin molecule containing sites for interaction with T cell receptors could lead to the formation of multimers. As will be appreciated, many methods can be used to obtain the coding region of the above affinity labels, including but not limited to, DNA cloning, DNA amplification methods. Some of the affinity labels and reagents for their detection are available commercially.

A preferred affinity label is a non-variable portion of an immunoglobulin molecule. Typically, such portions comprise at least a functional domain of the constant region of an immunoglobulin heavy chain using the carboxyl terminus of the Fc portion of a constant domain or the amino-terminal to the CH1 of the heavy or light chain. Suitable affinity labels may be obtained from IgG-1, -2, -3, or -4 subtypes, preferably IgG1. Preferably, a human immunoglobulin is used when the affinity label is intended for *in vivo* use for humans. Many DNA encoding immunoglobulin heavy chain constant regions is known or readily available from the literature. For example, Adams *et al.*, Biochemistry, 1980, 19:2711-2719; Gough *et al.*, Biochemistry, 19:2702-2710; Dolby *et al.*, 1980, Proc. Natl. Acad. Sci. U.S.A., 77:6031; Rice *et al.*, 1982, Proc. Natl. Acad. Sci. U.S.A., 79:7862-7867; Nature, 298:286-288; and Morrison *et al.*, 1984, Ann. Rev. Immunol. 2:241-264. Many immunological reagents and labeling systems are available for immunoglobulins, the  $\alpha 2M$  polypeptide-Ig fusion protein can be easily quantified by a variety of immunological techniques known in the art, such as enzyme-linked immunosorbent assay (ELISA), immunoprecipitation, flow cell sorting (FACS), *etc.* Similarly, if the affinity label is an epitope

antibodies, such reagents can be used with the techniques mentioned above to quantitate, and isolate the  $\alpha 2M$  polypeptide containing the affinity label. There is no need to develop specific antibodies to the  $\alpha 2M$  polypeptide.

A particularly preferred embodiment is a fusion of the hinge, the CH2 and CH3 domains of human immunoglobulin *al.*, 1996, J. Immunol. 156:442-49). This hinge region contains the cysteines which are normally involved in disulfide bonding with other cysteines. Since none of the cysteines are required for the peptide to function, these cysteine residues may optionally be substituted by another amino acid, for example, serine.

Various leader sequences known in the art can be used for the secretion of  $\alpha 2M$  polypeptide from bacterial and mammalian cells (Mol. Biol. 184:99-105). Leader peptides are selected based on their source, which may include bacterial, yeast, viral, animal, and mammalian sequences. The herpes virus glycoprotein D leader peptide is suitable for use in mammalian cells. A preferred leader peptide for use in mammalian cells can be the C region of the mouse immunoglobulin kappa chain (Bernard *et al.*, 1981, J. Biol. Chem. 256:5812-5816). Preferred leader sequences for targeting  $\alpha 2M$  in bacterial cells include, but are not limited to, the leader sequences of OmpA (Hobom *et al.*, 1995, Dev. Biol. Stand. 84:255-262), PhoA (Natl. Acad. Sci. 82:7212-16), OmpT (Johnson *et al.*, 1996, Protein Expr. Purif. 10:1-10), LamB and OmpF (Hoffman & Wright, 1985, Proc. Natl. Acad. Sci. USA 82:2149-54), *etc.* (Kadonaga *et al.*, 1984, J. Biol. Chem. 259:2149-54), and the *Staphylococcus aureus* A (Abrahmsen *et al.*, 1986, Nucleic Acids Res. 14:7487-7500), an endoglucanase (Lo *et al.*, Appl. Environ. Microbiol. 54:2287-2292), and synthetic signal sequences (MacIntyre *et al.*, 1990, Mol. Gen. Genet. 235:312-317).

DNA sequences encoding a desired affinity label can be readily obtained from libraries, produced synthetically, or from commercial suppliers, are suitable for the practice of this invention known in the art.

#### 4.5.3. Expression Systems

Nucleotide sequences encoding an hsp or  $\alpha 2M$  and an hsp-antigenic molecule or  $\alpha 2M$ -antigenic molecule fusion can be used in the practice of this invention.

expression vector to produce an expression construct for propagation in recombinant cells. An expression construct, as used herein, refers to a construct encoding an hsp,  $\alpha$ 2M, and/or antigenic molecule operably associated with regulatory regions which allows expression of the hsp,  $\alpha$ 2M and/or appropriate host cell. "Operably-associated" refers to an association in which the regulatory regions and the hsp,  $\alpha$ 2M and/or antigenic molecule polypeptide are joined and positioned in such a way as to permit transcription and translation of the hsp,  $\alpha$ 2M and/or antigenic molecule sequence. Such vectors may be used for the expression of hsps,  $\alpha$ 2M and/or antigens, but not limited to, plasmids, cosmids, phage, phagemids, or modified vectors include bacteriophages such as lambda derivatives, or plasmids or plasmid derivatives or the Bluescript vector (Stratagene). Typical vectors comprise a functional origin of replication for propagation in an appropriate host cell, one or more restriction endonuclease sites for insertion of a sequence or sequence encoding an antigenic molecule, and one or

For expression of hsps,  $\alpha$ 2M and/or antigenic molecules in host cells, a variety of regulatory regions can be used, for example, the CMV promoter, the cytomegalovirus (CMV) immediate early promoter, the RSV long terminal repeat (RSV-LTR) promoter. Inducible promoters in mammalian cells include but are not limited to those associated with the glucocorticoid responsive gene, mouse mammary tumor virus glucocorticoid responsive promoter (MMTV-LTR), the  $\beta$ -interferon gene, and the hsp70 gene (Williams *et al.*, 1985, *Res.* 49: 2735-42; Taylor *et al.*, 1990, *Mol. Cell. Biol.* 10: 165-7).

The following animal regulatory regions, which have been utilized in transgenic animals, can also be used for the expression of hsps and/or antigenic molecules in cells of a particular tissue type. The insulin gene control region which is active in pancreatic acinar cells (Swift *et al.*, 1984, *Cold Spring Harbor Symp. Quant. Biol.* 50: 399-409; Hanahan, 1985, *Nature* 315: 115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl *et al.*, 1984, *Cell* 38: 647-65; Alexander *et al.*, 1987, *Mol. Cell. Biol.* 7: 1533-538), mouse mammary tumor virus control region which is active in testicular cells (Leder *et al.*, 1986, *Cell* 45: 485-495), albumin gene control region which is active in liver (Pinkert *et al.*, 1987, *Genes Dev.* 1: 268-276),  $\alpha$ -fetoprotein gene control region which is active in liver (Krumlauf *et al.*, 1985, *Mol. Cell. Biol.* 5: 1315-1325).



Hammer *et al.*, 1987, Science 235: 53-58; alpha 1-antitrypsin gene active in the liver (Kelsey *et al.*, 1987, Genes Dev. 1: 161-171), beta region which is active in myeloid cells (Mogam *et al.*, 1985, Nature *et al.*, 1986, Cell 46: 89-94; myelin basic protein gene control region 5 oligodendrocyte cells in the brain (Readhead *et al.*, 1987, Cell 48: chain-2 gene control region which is active in skeletal muscle (Sar 286), and gonadotropic releasing hormone gene control region within hypothalamus (Mason *et al.*, 1986, Science 234: 1372-1378).

The efficiency of expression of the hsp,  $\alpha 2M$  or another cell may be enhanced by the inclusion of appropriate transcription expression vector, such as those found in SV40 virus, Hepatitis B immunoglobulin genes, metallothionein,  $\beta$ -actin (see Bittner *et al.* Enzymol. 153: 516-544; Gorman, 1990, Curr. Op. in Biotechnol. 1

The expression vector may also contain sequences for 15 and replication of the vector in more than one type of host cell, or into the host chromosome. Such sequences may include but are not origins, autonomously replicating sequences (ARS), centromere DNA. It may also be advantageous to use shuttle vectors that can be replicated at least two types of host cells.

20 In addition, the expression vector may contain selectable marker genes for initially isolating or identifying host cells that contain hsp,  $\alpha 2M$  and/or antigenic molecule. For long term, high yield production and/or antigenic molecules, stable expression in mammalian cells in selection systems may be used for mammalian cells, including, but 25 Herpes simplex virus thymidine kinase (Wigler *et al.*, 1977, Cell 1 guanine phosphoribosyltransferase (Szybalski and Szybalski, 1962 U.S.A. 48: 2026), and adenine phosphoribosyltransferase (Lowy *et al.* genes can be employed in *tk*-, *hgp*rt- or *ap*rt- cells, respectively. A resistance can be used as the basis of selection for dihydrofolate re 30 confers resistance to methotrexate (Wigler *et al.*, 1980, Natl. Acad. O'Hare *et al.*, 1981, Proc. Natl. Acad. Sci. U.S.A. 78: 1527); gpt, which confers resistance to mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl. Acad. neomycin phosphotransferase (neo), which confers resistance to tetracycline (Colberre-Garapin *et al.*, 1981, J. Mol. Biol. 150: 1); and hygromycin 35 (hyg), which confers resistance to hygromycin (Santerre *et al.*, 1981, J. Mol. Biol. 150: 1). Other selectable markers, such as but not limited to histidinol and Zeocin

In order to insert the hsp or  $\alpha 2M$  coding sequence an antigenic molecule into the cloning site of a vector, DNA sequence functions, such as promoters, must be attached to the respective c this, linkers or adapters providing the appropriate compatible restriction 5 to the ends of cDNA or synthetic DNA encoding an hsp,  $\alpha 2M$  or techniques well known in the art (Wu *et al.*, 1987, Methods Enzymol. Cleavage with a restriction enzyme can be followed by modification by digesting back or filling in single-stranded DNA termini before ligation. A desired restriction enzyme site can be introduced into a fragment 10 of the DNA by use of PCR with primers containing the desired restriction

The expression construct comprising an hsp,  $\alpha 2M$  molecule-coding sequence operably associated with regulatory elements is introduced into appropriate host cells for expression and production of antigenic molecule complexes of the invention without further cloning. 15 No. 5,580,859). The expression constructs may also contain DNA elements for integration of the coding sequence into the genome of the host cell by homologous recombination. In this instance, it is not necessary to employ an expression vector comprising a replication origin suitable for appropriate host cells to express the hsp,  $\alpha 2M$  and/or antigenic molecule in the host cells.

20 Expression constructs containing cloned hsp or  $\alpha 2M$  coding sequences for antigenic molecules can be introduced into host cells by a variety of techniques known in the art, including but not limited to: electroporation mediated transfection (Wigler *et al.*, 1977, Cell 11: 223-232), lipid-mediated transfection (Schaefer-Ridder *et al.*, 1982, Science 215: 166-168) 25 *et al.*, 1987, Proc. Natl. Acad. Sci. 84: 3344), and microinjection (Calkins *et al.*, 1987, J. Cell Biol. 100: 479-488).

For long-term, high-yield production of properly formed antigenic molecule complexes, stable expression in mammalian cells is preferred. Cells expressing hsp or  $\alpha 2M$  and antigenic molecules to produce hsp-peptide complexes 30 into the compositions of the present invention may be engineered to contain a selectable marker. By way of example but not limitation, the expression constructs, engineered cells may be grown for several days in an enriched media, and then are switched to a selective media. The selectable marker in the expression construct confers resistance to the selection agent. 35 cells to stably integrate the expression construct into their chromosomal DNA and to be expanded into cell lines. Such cells can be cultured

time while the hsp or  $\alpha 2M$  and antigenic molecule is expressed co

Any of the cloning and expression vectors describe synthesized and assembled from known DNA sequences by techn art. The regulatory regions and enhancer elements can be of a vari  
 5 natural and synthetic. Some vectors and host cells may be obtaine limiting examples of useful vectors are described in Appendix 5 o: Molecular Biology, 1988, ed. Ausubel *et al.*, Greene Publish. Assoc which is incorporated herein by reference; and the catalogs of com Clontech Laboratories, Stratagene Inc., and Invitrogen, Inc.

10 Alternatively, number of viral-based expression sys with mammalian cells for recombinant expression of hsps,  $\alpha 2M$  ar Vectors using DNA virus backbones have been derived from simia (Hamer *et al.*, 1979, Cell 17: 725), adenovirus (Van Doren *et al.*, 1 1653), adeno-associated virus (McLaughlin *et al.*, 1988, J. Virol. 6  
 15 papillomas virus (Zinn *et al.*, 1982, Proc. Natl. Acad. Sci. 79: 489). adenovirus is used as an expression vector, the donor DNA sequen adenovirus transcription/translation control region, *e.g.*, the late pr leader sequence. This chimeric gene may then be inserted in the a *vitro* or *in vivo* recombination. Insertion in a non-essential region  
 20 region E1 or E3) will result in a recombinant virus that is viable ar heterologous products in infected hosts (see, *e.g.*, Logan and Shen Sci. U.S.A. 81: 3655-3659).

Bovine papillomavirus (BPV) can infect many high man, and its DNA replicates as an episome. A number of shuttle v  
 25 developed for recombinant gene expression which exist as stable, copies/cell) extrachromosomal elements in mammalian cells. Typ contain a segment of BPV DNA (the entire genome or a 69% trans promoter with a broad host range, a polyadenylation signal, splice marker, and "poisonless" plasmid sequences that allow the vector t  
 30 *coli*. Following construction and amplification in bacteria, the exp transfected into cultured mammalian cells, for example, by the tech phosphate coprecipitation or electroporation. For those host cells t transformed phenotype, selection of transformants is achieved by t selectable marker, such as histidinol and G418 resistance. For exa  
 35 as pBCMGSNeo and pBCMGRHis may be used to express hsps,  $\alpha 2$  molecules (Karasuyama *et al.*, Eur. J. Immunol. 18: 97-104; Ohe *e*

Therapy 6: 325-33) which may then be transfected into a diverse range of cells for  $\alpha 2M$  or antigenic molecule expression.

Alternatively, the vaccinia 7.5K promoter may be used (Mackett *et al.*, 1982, Proc. Natl. Acad. Sci. U.S.A. 79: 7415-7419; Mackett *et al.*, 1982, Proc. Natl. Acad. Sci. U.S.A. 79: 857-864; Panicali *et al.*, 1982, Proc. Natl. Acad. Sci. U.S.A. 79: 49-54). If a human host cell is used, vectors based on the Epstein-Barr virus (EBV) nuclear antigen 1 (EBNA-1; a trans-acting replication factor) vectors can be used with a broad range of human host cells, e.g., El *et al.*, 1990, DNA Prot. Eng. Tech. 2: 14-18), pDR2 and  $\lambda$ DR2 (available from Pharmacia Laboratories).

Recombinant hsp,  $\alpha 2M$  and/or antigenic molecule expression can be achieved by a retrovirus-based expression system. In contrast to adenovirus, retrovirus can efficiently infect and transfer genes to a wide range of cell types including primary hematopoietic cells. In retroviruses such as Moloney murine leukemia virus, some of the viral gene sequences can be removed and replaced with an heterologous molecule coding sequence or a sequence encoding an antigenic molecule. Viral functions can be supplied in *trans*. The host range for infection can also be manipulated by the choice of envelope used for vector packaging.

For example, a retroviral vector can comprise a 5' long terminal repeat (LTR), a 3' LTR, a packaging signal, a bacterial origin of replication, and a selectable marker. The ND-associated antigenic peptide DNA is inserted into the 5' LTR and 3' LTR, such that transcription from the 5' LTR promoter produces the antigenic peptide DNA. The 5' LTR comprises a promoter, including but not limited to the U5 region, a U3 region and a primer binding site, in that order. Numerous LTR elements are well known in the art. A heterologous promoter or drug selection markers may also be included in the expression vector. (see McLaughlin *et al.*, 1990, Prog. Nucleic Acid Res. 41: 91-135; Morgenstern *et al.*, 1990, Nucleic Acid Res. 18: 3587-3596; J. Virol 70: 1792-1798; Boesen *et al.*, 1994, Biotherapy 6: 291-302; Gunzberg, 1993, Human Gene Therapy 4: 129-141; and Grossman *et al.*, 1993, Opin. in Genetics and Devel. 3: 110-114).

The recombinant cells may be cultured under standard conditions including temperature, incubation time, optical density, and media composition. The cells may be cultured under conditions emulating the nutritional and physiological conditions of a cell in which the hsp,  $\alpha 2M$  or antigenic molecule is endogenous. Standard culture conditions and media may be used to enhance production of the antigenic molecule.

or  $\alpha$ 2M-antigenic molecule complexes. For example, recombination under conditions that promote inducible hsp expression. Any test may be applied to establish the optimal conditions for producing  $\alpha$ 2M-antigenic molecule complexes.

5

#### 4.6. Therapeutic and Prophylactic Applications and Compositions of the Invention

In accordance with the present invention, a composition comprising an hsp or  $\alpha$ 2M, optionally an antigenic molecule, administered to a human subject with cancer, an infectious disease, or an autoimmune embodiment, "treatment" or "treating" refers to an amelioration of a disease, a neurodegenerative or amyloid disease or an autoimmune discernible symptom thereof. In another embodiment, "treatment" refers to amelioration of at least one measurable physical parameter associated with an infectious disease, a neurodegenerative or amyloid disease or an autoimmune symptom necessarily discernible by the subject. In yet another embodiment, "treatment" refers to inhibiting the progression of a cancer, an infectious disease, an amyloid disease, or an autoimmune disease, either physically, *e.g.*, stabilization of a physical symptom, physiologically, *e.g.*, stabilization of a physiological symptom. In yet another embodiment, "treatment" or "treating" refers to delaying the progression of a neurodegenerative or amyloid disease, or an autoimmune disease.

In certain embodiments, the compositions of the invention are administered to a human subject as a preventative measure against an infectious disease, a neurodegenerative or amyloid disease or an autoimmune disease. Herein, "prevention" or "preventing" refers to a reduction of the risk of developing a cancer, infectious disease, neurodegenerative or amyloid disease or an autoimmune disease. In one mode of the embodiment, the compositions of the invention are administered as a preventative measure to a human subject having a genetic predisposition to an infectious disease, neurodegenerative or amyloid disease or an autoimmune disease. In another mode of the embodiment, the compositions of the invention are administered as a preventative measure to a subject having a non-infectious disease, cancer, or to a subject facing exposure to an agent of an infectious disease.

##### 4.6.1. Target Infectious Diseases

Infectious diseases that can be treated or prevented by the present invention are caused by infectious agents including, but not limited to, bacteria, fungi, protozoa and parasites.

Viral diseases that can be treated or prevented by the present invention include, but are not limited to, those caused by hepatitis type C, influenza, varicella, adenovirus, herpes simplex virus type I (HSV-I), rinderpest, rhinovirus, echovirus, rotavirus, papilloma virus, papova virus, cytomegalovirus, echinovirus, coxsackie virus, mumps virus, measles virus, rubella virus, polio virus, immunodeficiency virus type I (HIV-I), and human immunodeficiency virus type II (HIV-II).

Bacterial diseases that can be treated or prevented by the present invention are caused by bacteria including, but not limited to, rickettsia, mycoplasma, neisseria and legionella.

Protozoal diseases that can be treated or prevented by the present invention are caused by protozoa including, but not limited to, kokzidia, and trypanosoma.

Parasitic diseases that can be treated or prevented by the present invention are caused by parasites including, but not limited to, rickettsia.

#### 4.6.2. Target Autoimmune Disease

Autoimmune diseases that can be treated by the present invention include, but are not limited to, insulin dependent diabetes, multiple sclerosis, systemic lupus erythematosus, scleroderma, polymyositis, chronic active hepatitis, primary biliary cirrhosis, pernicious anemia, autoimmune Addison's disease, vitiligo, gluten-sensitive enteropathy, Graves' disease, autoimmune neutropenia, idiopathic thrombocytopenia purpura, cirrhosis, pemphigus vulgaris, autoimmune infertility, Goodpasture's disease, pemphigoid, discoid lupus, ulcerative colitis, and dense deposit disease. The diseases referred to herein, include those exhibited by animal models of autoimmune diseases, such as, for example non-obese diabetic (NOD) mice for type 1 diabetes, autoimmune encephalomyelitis (EAE) mice for multiple sclerosis, and the like.

The methods of the present invention can be used to treat autoimmune diseases by reducing or eliminating the immune response to the self, or, alternatively, by reducing or eliminating a pre-existing autoimmune response to self tissues or organs transplanted to replace self tissues or organs damaged by disease.

#### 4.6.3. Target Cancers

Cancers that can be treated or prevented by the invention include, but are not limited to, human sarcomas and carcinomas, including fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteosarcoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, meningioma, melanoma, neuroblastoma, retinoblastoma; leukemia and acute myelocytic leukemia (myeloblastic, promyelocytic and erythroleukemia); chronic leukemia (chronic myelocytic leukemia and chronic lymphocytic leukemia); and polycythemia vera (Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, macroglobulinemia, and heavy chain disease. Specific examples are described in the sections below.

In a specific embodiment the cancer is metastatic. In another embodiment, the patient having a cancer is immunosuppressed by or has undergone anti-cancer therapy (e.g., chemotherapy radiation) prior to the use of the compositions of the invention alone or in combination with sensitizers.

##### 4.6.3.1 Colorectal Cancer Metastatic to the Liver

It has been estimated that approximately 226,600 Americans are diagnosed with cancers of the digestive tract in 2000. Most notably, the colon is the primary site for approximately 93,800 of these cases and the rectum is the site for another approximately 36,400 cases. Further, it is predicted that approximately 8,600 will die of colon cancer and another 8,600 will die of rectal cancer (Ca 2000, American Cancer Society (ACS), Atlanta, Georgia, 2000). Approximately 35% of patients who die of colon or rectal cancer have metastatic disease involving the tumors of the liver are from gastrointestinal primaries. Unfortunately,

the portal venous circulation to the liver, which represents the most common site of metastasis. The symptoms that lead patients with colorectal cancer to seek medical attention vary with the anatomical location of the lesion. For example, lesions of the sigmoid colon frequently ulcerate, which leads to chronic blood loss in the stool.

Patients with widespread metastasises have limited chemotherapy has little impact in this group of patients. In addition administered chemotherapy is often limited by the severity of toxic various agents, such as severe diarrhea, mucositis and/or myelosuppression techniques, including hepatic radiation, systemic chemotherapy, hepatic tumor embolization and immunotherapy have all been explored, but proven ineffectual in prolonging patient survival.

Accordingly, as an example of the method of the invention, the metal-organic  
molecule complexes are prepared from the cancer cells of a patient with  
colorectal cancer, with or without liver metastases, combined with  
35 and administered to the patient, via one of many different routes of  
administration, the preferred route being intradermally at different anatomical sites, e.g.



belly, right belly, left thigh, right thigh, *etc.* The site of injection varies with the site of weekly injection.

#### 4.6.3.2 Hepatocellular Carcinoma

- 5 Hepatocellular carcinoma is generally a disease of the United States. Although many factors may lead to hepatocellular carcinoma, it is limited to those persons with preexisting liver disease. Approximately 10 patients in the United States with hepatocellular carcinoma have a four percent of individuals with a cirrhotic liver eventually develop carcinoma (Niederhuber (ed.), 1993, *Current Therapy in Oncology*). The risk is highest in patients whose liver disease is caused by chronic viral or hepatic B viral infection (Bradbear *et al.*, 1985, *J. Natl. Cancer* 60:1129); hepatitis C virus infection has also emerged in the past decade (Colombo, 1999, Baillieres Best Pract Res Clin Gastroenterol). Other causes of cirrhosis that can lead to hepatocellular carcinoma include 15 and hepatic fibrosis caused by chronic administration of methotrexate. The symptoms of hepatocellular carcinoma are the development of a palpable mass in the upper quadrant or epigastrium, accompanied by weight loss. In patients with the development of hepatocellular carcinoma is preceded by ascites, peripheral 20 relatively abrupt clinical deterioration. In most cases, abnormal liver function tests such as serum aminotransferase and alkaline phosphatase. CT scans of the liver are used to determine the anatomical location of hepatocellular carcinoma and also provide orientation for percutaneous biopsy. Approximately 70 percent of patients with hepatocellular carcinoma have an elevated alpha-fetoprotein concentration (McIntire *et al.*, 1975, *Cancer Res* 35:1129). Alpha-fetoprotein concentration correlates with the extent of the disease.

Radical resection offers the only hope for cure in patients with hepatocellular carcinoma. Such operative procedures are associated with five-year survival rates of 30 percent. Liver transplantation may improve survival of some patients. However, most patients are not surgical candidates because of extensive tumor pattern or scarcity of compatible donor organs.

Chemotherapeutic agents have been administered either intravenously or through an intrahepatic arterial catheter. Such therapy has some effect when combined with irradiation to the liver. Reductions in the size of measurable tumors 35 have been reported in some patients treated with either systemic chemotherapy or intrahepatic chemotherapy. However, chemotherapy often induces immunosuppression.

the tumor to disappear completely and the duration of response is  
 patients with hepatocellular carcinoma is negatively correlated wi  
 to the lungs or bone. Median survival for patients is only four to  
 embodiment, the present invention provides compositions and me  
 5 specific immunity in individuals suffering from hepatocellular ca  
 the progression of the neoplastic disease and ultimately irradiate  
 neoplastic cells. Thus the present invention also provides compo  
 $\alpha$ 2M, saponins, and hepatocellular carcinoma-antigenic molecule  
 promoting immunity to the disease and for inhibiting cancer cell  
 10 metastasis. These compositions can be applied alone or in combi  
 with biological response modifiers.

#### 4.6.3.3 Breast Cancer

Another specific aspect of the invention relates to  
 15 cancer. The American Cancer Society estimated that in 2000, 18  
 will be diagnosed with breast cancer and 41,200 will succumb to  
 & Figures 2000, American Cancer Society (ACS), Atlanta, Georg  
 breast cancer the second major cause of cancer death in women, r  
 cancer. The treatment of breast cancer presently involves surgery  
 20 therapy and/or chemotherapy. Consideration of two breast cance  
 receptors and disease extent, has governed how hormonal therapi  
 chemotherapy are sequenced to improve survival and maintain or  
 wide range of multidrug regimens have been used as adjuvant the  
 patients, including, but not limited to combinations of 2 cyclopho  
 25 vincristine methotrexate, 5-fluorouracil and/or leucovorin. In a s  
 present invention provides compositions comprising hsps or  $\alpha$ 2M  
 cancer-antigenic molecules and methods for enhancing specific ir  
 and neoplastic mammary cells in women. The present invention  
 comprising hsps or  $\alpha$ 2M, saponins, and breast cancer-antigenic m  
 30 preventing the development of neoplastic cells in women at enhai  
 and for inhibiting cancer cell proliferation and metastasis. These  
 applied alone or in combination with each other or with biologica

#### 4.6.3.4 Autologous Antigenic Molecu

35 The specific immunogenicity of hsp or  $\alpha$ 2M -antig  
 derives from the antigenic molecules therein. In a preferred emb

compositions of the invention comprising autologous antigenic m  
 cancer immunotherapy. In one embodiment, the composition of t  
 complexes of antigenic molecules and hsps purified from cancero  
 individual to whom they are to be administered. In other embodi  
 5 molecules are dissociated from the hsp-antigenic molecule compl  
 cancerous cells and incorporated into the compositions of the inv  
 covalently or non-covalently complexed to  $\alpha$ 2M, covalently comp  
 QS-21, or covalently or non-covalently complexed to another hsp  
 recombinant hsp. This approach offers the advantage of using an  
 10 to cancers that are potentially antigenically distinct. Additionally  
 embodiments, cancer immunotherapy does not depend on the ava  
 CTLs nor does it require definition of the antigenic epitopes of ca

#### 4.6.4. Target Neurodegenerative

15 Neurodegenerative diseases that can be treated or p  
 the present invention include, but are not limited to, Alzheimer's  
 cognitive function, senile dementia, Parkinson's disease, amyotro  
 Wilson's Disease, cerebral palsy, progressive supranuclear palsy,  
 dementia, prion diseases, spongiform encephalopathies, Creutzfe  
 20 polyglutamine diseases, Huntington's disease, myotonic dystroph  
 ataxia, Gilles de la Tourette's syndrome, seizure disorders, epilep  
 disorder, stroke, brain trauma, spinal cord trauma, AIDS dementi  
 retinal ischemia, glaucoma, autonomic function disorder, hyperten  
 disorder, schizophrenia, or schizoaffective disorder.

25 Amyloid diseases that can be treated or prevent by  
 present invention include but are not limited to are diseases chara  
 extracellular deposition of protein and/or peptide fibrils which fo  
 plaques, including but not limited to type II diabetes and amyloid  
 chronic inflammatory or infectious disease states and malignant  
 30 Certain amyloid disease such as Alzheimer's disease and prion di  
 Jacob disease, are neurodegenerative diseases.

#### 4.7. Formation of Covalent Complexes

In certain embodiments of the present invention,  $\alpha$   
 35 covalently complexed to hsps,  $\alpha$ 2M and/or saponin. The covalen  
 molecules to hsps,  $\alpha$ 2M and/or saponin is done prior to administ

compositions of the invention are formulated. Wherein covalent endogenous hsp-peptide complex is desired, the complex is preferred purification from cells or tissues. In one embodiment, antigenic molecules are coupled to hsps,  $\alpha$ 2M and/or saponin by chemical crosslinking. (5 methods are well known in the art. For example, in a preferred embodiment, crosslinking may be used. Glutaraldehyde crosslinking has been used to form covalent complexes of peptides and hsps (see Barrios *et al.*, 1992, 1365-1372). Preferably, 1-2 mg of complex is crosslinked in the glutaraldehyde for 2 hours. Glutaraldehyde is removed by dialysis 10 buffered saline (PBS) overnight (Lussow *et al.*, 1991, Eur. J. Immunol. 21: 1365-1372).

In another embodiment, the antigenic molecules and saponin are crosslinked by ultraviolet (UV) crosslinking.

For  $\alpha$ 2M-antigenic molecule complexes, covalent complexes are prepared by preparing the complexes according to the method described by Giedt *et al.*, 1991, J. Biol. Chem. 266: 6009-6014. 15 Biochemistry, 37: 6009-6014.

#### 4.8. Combination With Adoptive Immunotherapy

Adoptive immunotherapy refers to a therapeutic approach for cancer or infectious diseases in which immune cells are administered to a patient. (20 cells mediate either directly or indirectly specific immunity to tumor antigens or components thereof or regression of the tumor or treatment of infection. In one case may be (see U.S. Application No. 08/527,546, filed September 16, 1996, Patent No. 5,985,270 issued November 16, 1999, which is incorporated by reference in its entirety). As an optional step, in accordance with the method of the invention, 25 are sensitized with hsps or  $\alpha$ 2M complexed with antigenic (or immunogenic) molecules used in adoptive immunotherapy. See Binder *et al.*, 2000, Nature Medicine 6: 154.

In a specific embodiment, therapy by administration of compositions comprising an hsp or  $\alpha$ 2M, an antigenic molecule, and an immunogenic molecule, 30 desired route of administration, may optionally be combined with administration of APC sensitized with hsp- or  $\alpha$ 2M- antigenic molecule complex. In Section 4.10 herein, the hsp- or  $\alpha$ 2M- peptide complex-sensitized APC is administered alone, in combination claimed compositions, or before or after administration of the claimed compositions. Furthermore, the mode of administration is 35 but not limited to, e.g., mucosally or intramuscularly, although intravenous or subcutaneously is preferred.

**4.8.1. Obtaining Antigen-Presen**

The APCs, including but not limited to macrophage cells, are preferably obtained by production *in vitro* from stem and human peripheral blood or bone marrow as described by Inaba *et al.*

5 176:1693-1702.

APCs can be obtained by any of various methods. In a preferred aspect human macrophages are used, obtained from human peripheral blood. For example but not limitation, macrophages and dendritic cells can be used.

Mononuclear cells are isolated from peripheral blood of the patient to be treated), by Ficoll-Hypaque gradient centrifugation. The cells are incubated in tissue culture dishes which are pre-coated with the patient's own human serum. The cells are incubated at 37°C for 1 hour, then non-adherent cells are removed by pipetting. To the adherent cells left in the dish, is added EDTA in phosphate-buffered saline and the dishes are left at room temperature for 15 minutes. The cells are harvested, washed with RPMI buffer and resuspended in RPMI medium. Increased numbers of macrophages may be obtained by incubating with macrophage-colony stimulating factor (M-CSF); increased numbers may be obtained by incubating with granulocyte-macrophage-colony stimulating factor (GM-CSF) as described in detail by Inaba *et al.*, 1992, J. Exp. Med. 176:1693-1702.

20

**4.8.2. Sensitization of Antigen Presenting Cells with hsp- or  $\alpha$ 2M- Antigenic Complexes**

APCs are sensitized with complexes of hsp or  $\alpha$ 2M and antigen. This is preferably achieved by incubating the cells *in vitro* with the complexes. Sensitization is achieved by incubating the APCs *in vitro* with the hsp- or  $\alpha$ 2M-antigen complexes at 37°C for 15 minutes to 24 hours. By way of example,  $4 \times 10^7$  macrophages can be incubated with 10 microgram gp96-peptide complexes or 100 microgram hsp90-peptide complexes per ml at 37°C for 15 minutes in plain RPMI medium. The cells are washed three times and resuspended in sterile physiological medium, at a convenient concentration (e.g.,  $1 \times 10^6$  cells/ml) for use in a patient. Preferably, the patient into which the sensitized APCs are injected is the same from which the APCs were originally isolated (autologous embryo).

30

Optionally, the ability of sensitized APCs to stimulate antigen-specific, class I-restricted cytotoxic T-lymphocytes (CTLs) is tested. Alternatively, their ability to stimulate CTLs to release tumor necrosis factor, or other cytokines, or to kill targets of such CTLs.

35

**4.8.3. Reinfusion of Sensitized A**

In one embodiment, APCs sensitized with complete molecule complexes or  $\alpha 2M$ -antigenic molecule complexes are administered systemically, preferably intravenously, by conventional clinical procedures. Patients generally receive from about  $10^6$  to about  $10^{12}$  sensitized APCs, depending on the condition of the patient. In some regimens, patients may optionally receive a dosage of a biological response modifier including but not limited to IFN- $\gamma$ , IL-2, IL-4, IL-6, TNF or other cytokine growth factor.

**4.9. Passive Immunotherapy**

The hsp or  $\alpha 2M$ / antigenic molecule/ saponin conjugate can also be used for passive immunotherapy against cancers and infectious diseases. Passive immunity is the short-term protection of a host, achieved by the administration of pre-formed antibody directed against a heterologous organism. The invention comprising antigenic molecules that display the epitopes of an infectious organism may be used to elicit an immune response. The response may be removed from the subject and used for treatment or prevention of the infectious organism in another subject.

**4.10. Formulation, Administration and**

The hsp or  $\alpha 2M$ / antigenic molecule/ saponin conjugate of the invention may be formulated into pharmaceutical preparations for use in mammals, preferably humans, for treatment or prevention of cancer or infectious diseases. Compositions comprising a compound of the invention in a compatible pharmaceutical carrier may be prepared, packaged and used for treatment of the indicated tumor(s), infectious disease or autoimmune disease.

Drug solubility and the site of absorption are factors that are considered when choosing the route of administration of a therapeutic embodiment of the invention, the present compositions may be administered by a desired route of administration, and preferably intradermally or subcutaneously. Advantages of intradermal or subcutaneous administration include rapid absorption, respectively. As an alternative to intradermal or subcutaneous administration, the compositions of the invention can be administered by other routes of administration include, but are not limited to, oral, rectal, intramuscular, intravenous, and intranasal. Preparations for mucosal administrations are suitable in various forms and are described below.

Water-soluble compositions of the invention may appropriate buffer, for example, phosphate buffered saline or other compatible solutions, preferably sterile. Alternatively, if the compound is soluble in aqueous solvents, then it may be formulated with a non-ionic surfactant such as polyethylene glycol. The compositions of the invention may be in the form of a powder or from a saponin stock aqueous solution by simple admixture with antigenic molecules. The compositions may be in any form, particularly when recombinant forms of these molecules are used as compounds and their physiologically acceptable solvates may be administered by inhalation or insufflation (either through the mouth, buccal, parenteral, or rectal administration or, in the case of a solid tumor.

For oral administration, the pharmaceutical preparation may be in the form, for example, solutions, syrups or suspensions, or may be in the form of a powder for reconstitution with water or other suitable vehicle before use. The preparations may be prepared by conventional means with pharmaceutically acceptable suspending agents (*e.g.*, sorbitol syrup, cellulose derivatives or hydroxypropyl methylcellulose); emulsifying agents (*e.g.*, lecithin or acacia); non-aqueous vehicles (*e.g.*, esters, or fractionated vegetable oils); and preservatives (*e.g.*, methylparaben, hydroxybenzoates or sorbic acid). The pharmaceutical compositions may be in the form, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.*, pregelatinized starch, hydroxypropyl methylcellulose); fillers (*e.g.*, lactose, cellulose or calcium hydrogen phosphate); lubricants (*e.g.*, magnesium stearate, silica); disintegrants (*e.g.*, potato starch or sodium starch glycolate); and surfactants (*e.g.*, sodium lauryl sulphate). The tablets may be coated by methods

Preparations for oral administration may be suitable for controlled release of the active compound.

For buccal administration, the compositions may be in the form of lozenges formulated in conventional manner.

The compounds may be formulated for parenteral injection, *e.g.*, by bolus injection or continuous infusion. Formulations may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers with a suitable preservative. The compositions may take such forms as suspensions, emulsions, or solutions in oily or aqueous vehicles, and may contain formulatory agents such as stabilizing and/or dispersing agents. Alternatively, the active ingredient

form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen:

The compounds may also be formulated in rectal or vaginal suppositories or retention enemas, *e.g.*, containing conventional solid carriers such as cocoa butter or other glycerides.

5 In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compositions may contain suitable polymeric or hydrophobic materials (for example, as an emulsion or as a depot oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a soluble salt. Liposomes and emulsions are well known examples of suitable carriers for hydrophilic drugs.

For administration by inhalation, the compositions of the present invention are conveniently delivered in the form of an aerosol or suspension from pressurized packs or a nebulizer, with the use of a suitable propellant such as dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane or other suitable gas. In the case of a pressurized aerosol the dosage may be determined by providing a valve to deliver a metered amount. Carriers such as *e.g.*, gelatin for use in an inhaler or insufflator may be formulated in a solid form of the compound and a suitable powder base such as lactose or starch.

The compositions may, if desired, be presented in a package or container which may contain one or more unit dosage forms containing the compound. The package may for example comprise metal or plastic foil, such as a blister pack. The package or dispenser device may be accompanied by instructions for administration.

25 The invention also provides kits for carrying out the invention. Such kits comprise in one or more containers therapeutically or prophylactically effective amounts of the hsp or  $\alpha 2M$ , uncomplexed or covalently complexed to an antigenic molecule, and a saponin. Kits for the treatment or prevention of cancer or infectious disease comprise at least one of the following: (a) uncomplexed or complexed to hsp or  $\alpha 2M$ ; and a saponin, uncomplexed or complexed to an antigenic molecule. Each kit component is preferably purified and in an acceptable form. Kits directed to the treatment or prevention of cancer optionally further comprise in a separate container, APCs, which are used to sensitize APCs. If the APCs are not sensitized, the kit may further comprise a purified antigenic molecule for sensitizing the APCs. The APCs are preferably purified. Each kit component is preferably lyophilized or dessicated (*i.e.*, in powder form), or may be in the form of a solution or suspension.



acceptable solution, *e.g.*, in combination with sterile saline, dextran solution, or other pharmaceutically acceptable sterile fluid. When the active component is lyophilized or desiccated, the kit optionally further includes a pharmaceutically acceptable solution (*e.g.*, saline, dextrose solution) to reconstitute the material to form a solution for injection purposes.

In another embodiment, a kit of the invention further includes a syringe, preferably packaged in sterile form, for injecting the composition, a packaged alcohol pad. Instructions are optionally included for administration of the compositions of the invention by a clinician or by the patient.

#### 4.11. Monitoring of Effects During Cancer Immunotherapy with the Compositions

The effect of immunotherapy with the hsp or  $\alpha 2M$  saponin compositions of the present invention on development of neoplastic diseases can be monitored by any methods known to one of skill in the art, including but not limited to measuring: a) delayed hypersensitivity; b) cellular immunity; c) activity of cytolytic T-lymphocytes *in vitro*; d) antigenic molecules, *e.g.*, carcinoembryonic (CEA) antigens; e) levels of tumors using techniques such as a computed tomographic (CT) scan; f) levels of putative biomarkers of risk for a particular cancer in individuals; g) changes in the morphology of tumors using a sonogram.

##### 4.11.1. Delayed Hypersensitivity

Delayed hypersensitivity skin tests are of great value in assessing immunocompetence and cellular immunity to an antigen. Inability to respond to common skin antigens is termed anergy (Sato *et al.*, 1995, *Clin. Immunol.* 77: 43).

Proper technique of skin testing requires that the antigen be stored at 4°C, protected from light and reconstituted shortly before use. A small volume ensures intradermal, rather than subcutaneous, administration of the antigen. 48 hours after intradermal administration of the antigen, the largest wheal, erythema and induration are measured with a ruler. Hypoactivity of a group of antigens is confirmed by testing with higher concentrations. In ambiguous circumstances, by a repeat test with an intermediate concentration.

##### 4.11.2. Activity of Cytolytic T-Lymphocytes

$8 \times 10^6$  peripheral blood derived T lymphocytes isolated

Hypaque centrifugation gradient technique, are restimulated with treated tumor cells in 3ml RPMI medium containing 10% fetal calf serum. In some experiments, 33% secondary mixed lymphocyte culture supernatant is added to the culture medium as a source of T cell growth factors.

- 5 In order to measure the primary response of cytolytic T cells to tumor immunization, T cells are cultured without the stimulator tumor cells. After six days, T cells are restimulated with antigenically distinct cells. After six days, the cells are tested for cytotoxicity in a 4 hour  $^{51}\text{Cr}$ -release assay. The spontaneous release of targets should reach a level less than 20%. For the anti-MHC class II assay, a tenfold concentrated supernatant of W6/32 hybridoma is added to the assay at a concentration of 12.5% (Heike *et al.*, J. Immunotherapy 15:165-175, 1992).

#### 4.11.3. Levels of Tumor Antigen

- Although it may not be possible to detect unique tumor antigens, many tumors display antigens that distinguish them from normal cells. Monoclonal antibody reagents have permitted the isolation and characterization of these antigens and have been invaluable diagnostically for distinguishing transformed cells and for definition of the cell lineage of transformed cells. Characterized human tumor-associated antigens are the oncofetal antigens. These are expressed during embryogenesis, but are absent or very difficult to detect in adult tissue. The prototype antigen is carcinoembryonic antigen (CEA), which is found on fetal gut and human colon cancer cells, but not on normal cells. CEA is shed from colon carcinoma cells and found in the serum of cancer patients. The presence of this antigen in the serum could be used to screen for cancer. However, patients with other tumors, such as pancreatic cancer, have elevated serum levels of CEA. Therefore, monitoring the level of CEA in cancer patients undergoing therapy has proven useful for predicting responses to treatment.

- Several other oncofetal antigens have been useful for monitoring human tumors, e.g., alpha-fetoprotein, an alpha-globulin. Alpha-fetoprotein, which is found in the serum of patients with liver tumors and yolk sac cells, is found in the serum of patients with liver tumors and can be used as a marker of disease status.

#### 4.11.4. Computed Tomographic

- CT remains the choice of techniques for the accurate diagnosis of tumors. CT has proved more sensitive and specific than any other imaging technique.

of metastases.

#### 4.11.5. Measurement of Putative

The levels of a putative biomarker for risk of a spe  
5 to monitor the effect of hsp bound to peptide complexes. For exam  
enhanced risk for prostate cancer, serum prostate-specific antigen  
measured by the procedure described by Brawer *et al.*, 1992, J. Ur  
Catalona *et al.*, 1993, JAMA 270:948-958; or in individuals at risk  
CEA is measured by techniques well known in the art; and in indi  
10 for breast cancer, 16- $\alpha$ -hydroxylation of estradiol is measured by t  
by Schneider *et al.*, 1982, Proc. Natl. Acad. Sci. USA 79:3047-30.

#### 4.11.6. Sonogram

A sonogram remains an alternative choice of techn  
15 staging of cancers.

#### 4.12. Combination Therapy for Cancer

The present hsp or  $\alpha$ 2M/ antigenic molecule/ sapor  
administered together with treatment with irradiation or one or mo  
20 agents. For irradiation treatment, the irradiation can be gamma ray  
general overview of radiation therapy, see Hellman, Chapter 12: I  
Therapy Cancer, in: Principles and Practice of Oncology, DeVita  
Lippencott Company, Philadelphia. Useful chemotherapeutic age  
taxol, mercaptopurine, thioguanine, hydroxyurea, cytarabine, cycl  
25 ifosfamide, nitrosoureas, cisplatin, carboplatin, mitomycin, dacart  
etoposides, campathecins, bleomycin, doxorubicin, idarubicin, da  
plicamycin, mitoxantrone, asparaginase, vinblastine, vincristine, v  
docetaxel. In a specific embodiment, a composition of the invention  
concurrently with radiation therapy or one or more chemotherapeu  
30 specific embodiment, chemotherapy or radiation therapy is admin  
to administration of a present composition, preferably at least an h  
a day, a week, a month, more preferably several months (e.g., up t  
subsequent to administration of a composition of the invention.

35

#### 4.13. Dosages

In the dosage regimens recited below, an hsp or  $\alpha$ 2

without complexed covalently or non-covalently bound antigenic molecule administered together with a saponin. Preferred routes of administration are intradermal or subcutaneous, wherein the sites of administration are varied. For example, the dosage regimens may be given once weekly for 4 to 5 weeks, and the mode of administration may be varied with each administration. Each site of administration may be varied sequentially. Thus, by way of limitation, the first injection can be given, either intradermally or subcutaneously, in the left arm, the second on the right arm, the third on the left belly, the fourth on the right thigh, the fifth on the left thigh, the sixth on the right thigh, *etc.* The same or different sites may be used after a gap of one or more injections. Also, split injections can be given. For example, half the dose can be given in one site and the other half in another site on the same day.

After 4-6 weeks, further injections are preferably given at intervals of one month over a period of time of one month. Later injections can be given at intervals of 15 days. The frequency of later injections can be modified depending upon the patient's clinical responsiveness to the immunotherapy. Alternatively, the mode of administration can be sequentially varied, e.g., weekly administrations are given in sequence intradermally, subcutaneously, and intramuscularly.

The saponin or  $\alpha 2M$  and hsp or  $\alpha 2M$ -antigenic molecule composition of the invention may be administered to mammalian subjects, e.g., primate subjects, horses, cows, pigs, *etc.*, and preferably humans.

An amount of saponin from about 0.1  $\mu g$  to about 1000  $\mu g$  per hsp or  $\alpha 2M$ /antigenic molecule/saponin composition of the invention can be anywhere within the range of 3, 5, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 110, 125, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000  $\mu g$ . Saponin may be used. The most preferred saponin dose is generally about 100  $\mu g$ .

It will be understood that, where reference is made to an antigenic molecule as a component of a composition, an antigenic peptide or full-length antigenic molecule (e.g. having more than 50 amino acid residues). The amount of an antigenic molecule when administered as a complex, is determined by the amount which is complexed non-covalently to an hsp or  $\alpha 2M$ . The amount of an antigenic molecule as free antigenic molecule together in a composition with an hsp or  $\alpha 2M$  administered in the range of from 10 to 1500  $\mu g$ , with from 25 to 1000  $\mu g$ , e.g., 10, 30, 50, 75, 100, 300, 500, 1000 or 1500  $\mu g$  may be used.

In certain embodiments, a dose of the composition comprises an amount of hsp or  $\alpha 2M$ /antigenic molecule/saponin composition of the invention.

or more other components of the composition is not therapeutic as when the components are combined. Accordingly, an immunogen invention comprises can sub-immunogenic amounts of its individual components, for example, an hsp-antigenic molecule in an amount t  
 5 immunogenic or sub-therapeutic, but is immunogenic or therapeutic saponin according to the disclosed methods. Methods of assaying the individual components or subcombinations of the composition described in Section 4.4, *supra*.

#### 10 4.13.1. Autoimmune Diseases

For the prevention or treatment of an autoimmune  
 dosages of the hsps or  $\alpha$ 2M in the compositions of the invention a  
 fold lower than those disclosed in U.S. Patent No. 6,007,821, which  
 by reference. For example, 0.1  $\mu$ g to 1 mg of an hsp or  $\alpha$ 2M may  
 15 preferred and 1-10  $\mu$ g most preferred. The hsp or  $\alpha$ 2M component  
 anywhere within the 0.1  $\mu$ g to 1 mg range; *e.g.*, 0.2, 0.3, 0.5, 1, 3,  
 1000  $\mu$ g may be used. Preferred routes of administration are intra  
 with lower dosages preferred for the former and 5 to 10 fold higher  
 the latter.

20 Additional dosages of the hsps or  $\alpha$ 2M in the composition  
 range from about 0.05  $\mu$ g to about 5000  $\mu$ g, preferably from about  
 In mammals, a range of from about 5  $\mu$ g to about 500  $\mu$ g, either in  
 subcutaneously, is preferred, with from about 5  $\mu$ g to about 200  $\mu$   
 from about 0.5  $\mu$ g to about 100  $\mu$ g intradermally being most preferred

#### 25 4.13.2. Cancer and Infectious Diseases

For the prevention or treatment of cancer or infectious  
 dosages of the the hsps or  $\alpha$ 2M components of the compositions are  
 generally similar to those disclosed in U.S. Patent No. 6,017,540,  
 30 herein by reference. For example, 0.01  $\mu$ g to 1 mg hsps or  $\alpha$ 2M may  
 preferred. The non-saponin component can be anywhere within the  
*e.g.*, 0.02, 0.05, 0.1, 0.2, 0.5, 1, 2, 5, 10, 20, 50, 100, 200, 500 or  
 be used. Preferred routes of administration are intradermal and sub  
 dosages preferred for the former and 5 to 10 fold higher dosages preferred

35

**4.13.3. Neurodegenerative and**

For the prevention or treatment of neurodegenerative diseases, the preferred dosages of the hsp or  $\alpha 2M$  components of the composition (optionally complexed to neurodegenerative disease- or amyloid antigenic molecules) are generally similar to those disclosed in U.S. Pat. Nos. 09/489,215 and 09/489,216, each of which is incorporated by reference in its entirety, although the dosages may be up to 10-fold less than the foregoing U.S. applications. Generally, the dosage is in the range of 5000  $\mu\text{g}$  for a human patient, the preferred human dosage being 1000  $\mu\text{g}$  for a mouse. In specific embodiments, the non-saponin component of the composition is in the 0.01  $\mu\text{g}$  to 1 mg range; e.g., 0.02, 0.05, 0.1, 0.2, 0.5, 1, 2, 5, 10, 100, 1000  $\mu\text{g}$  hsp or  $\alpha 2M$ , preferably complexed to a neurodegenerative disease-associated antigenic molecule, may be used.

Compositions of the invention comprising the dosages disclosed herein are preferably given once weekly for a period of about 4-6 weeks, and the administration is preferably varied with each administration.

In a specific embodiment, the compositions of the invention are used for the prevention or treatment of a neurodegenerative disease by administration to the cerebrospinal fluid (CSF) of a patient by means of injection. In another embodiment, administration is in or near the lesioned area of the brain.

In a preferred example, intradermal administration of the composition is varied sequentially. Thus, by way of example, the first injection may be given subcutaneously on the left arm, the second on the left belly, the fourth on the right belly, the fifth on the left thigh, etc. The same site may be repeated after a gap of one or more days. Multiple injections may be given. Thus, for example, half the dose may be given at one site and the other half on another site on the same day.

Alternatively, the mode of administration is sequential. Thus, the injections are given in sequence intradermally, intramuscularly, or intraperitoneally.

After 4-6 weeks, further injections are preferably given over a period of time of one month. Later injections may be given at longer intervals. Later injections may be modified, depending upon the patient's clinical responsiveness to the immunotherapy.

#### 4.14. Methods of Making Pharmaceutical Comprising an hsp or $\alpha$ 2M, an Antigenic Molecule and a Saponin

The present invention encompasses methods of making pharmaceutical compositions comprising an hsp or  $\alpha$ 2M, an antigenic molecule the antigenicity of an antigen associated cancer, infectious disease, ne amyloid disease or autoimmune disease, and a saponin adjuvant, c saponin, the antigenic molecule and the hsp or  $\alpha$ 2M under condit pharmaceutical composition.

In certain embodiments, saponin is combined with and then the resulting mixture combined with the hsp or  $\alpha$ 2M. Th complexed molecules or complexed to another antigenic molecule saponin and the antigenic molecule are subjected to conditions th covalent saponin-antigenic molecule complex prior to combining or  $\alpha$ 2M.

In an embodiment where the saponin is combined molecule prior to combining the mixture with the hsp or  $\alpha$ 2M, the complexed to any other molecule. In an alternative embodiment, t complexed to another antigenic molecule. The hsp or  $\alpha$ 2M can bi covalently complexed to the second antigenic molecule. In yet an hsp or  $\alpha$ 2M is in the form of a fusion protein comprising the hsp c antigenic molecule.

Alternatively, the hsp or  $\alpha$ 2M can be combined with and then the resulting mixture can be combined with the saponin. antigenic molecule is combined with hsp or  $\alpha$ 2M under condition of a complex comprising the antigenic molecule and the hsp or  $\alpha$ 2M and the antigenic molecule are combined prior to combining saponin, the saponin is optionally covalently attached to another a

In an alternative embodiment, the hsp or  $\alpha$ 2M, the molecule are combined simultaneously. Optionally, the resulting conditions that promote covalent binding between the antigenic m  $\alpha$ 2M and/or between the antigenic molecule and the saponin.

For pharmaceutical compositions prepared by mix antigenic molecule complex with a saponin, the hsp- or  $\alpha$ 2M-anti can be prepared *in vitro* or purified from a cell or tissue, as descri The cell or tissue is optionally transfected with a nucleic acid enc

## 5. Example: Administration of an hsp or $\alpha$ 2M, and Saponin in Two UV-Induced Mouse Ca

Two UV-induced carcinomas are available in C3H compositions and methods of the invention: (i) the highly-immunogenic 6138 and (ii) the less immunogenic 6139SJ carcinoma (Ward *et al.*, 1981).

For example, an hsp such as gp96 is prepared from 6138 and 6139SJ carcinomas, or  $\alpha$ 2M is prepared from 6138 and 6139SJ carcinoma: expression construct, by the procedures described in Section 4 above. The antigenic molecules derived therefrom, are then administered as an adjuvant.

### 5.1. Prevention Modality Using gp96

Materials and Methods. The ability of hsp or  $\alpha$ 2M saponin compositions to prevent development of UV-induced carcinoma follows. Groups of female C<sub>3</sub>H/HeN mice (obtained from the National Cancer Institute, Frederick, MD), weighing approximately 25g each, are used in the experiments below.

Groups of mice are given twice at a ten day interval: (i) buffer saline (PBS), (ii) 0.1, 1, 10, 25, 50, or 100  $\mu$ g/mouse of, for complexes derived from UV6138 carcinomas, or (iii) 0.1, 1, 10, 25, 50, or 100  $\mu$ g of gp96-peptide complexes derived from UV6139SJ carcinoma. For each dosage series, 0, 1.6, 10, 20, 50, or 100  $\mu$ g of saponin fraction (from lyophilized powder) is mixed with the gp96-peptide complex and coadministered. Control sets of mice receive the dosage series of PBS.

Mice are challenged with  $10^7$  cells from either the UV6138 or UV6139SJ carcinoma 15 days after the second injection. Tumors are observed at 30 day intervals. Since the UV6138 tumor is a regressor tumor, mice are observed for 60 days after the second injection with PBS or gp96 in order to permit tumor regression. The UV6139SJ challenged mice are not irradiated.

Results. It has previously been shown that administration of gp96 from the UV6138 carcinoma rendered the mice immune to the UV6138 challenge (see U.S. Patent No. 5,837,251). Conversely, gp96 isolated from the UV6139SJ conferred resistance to the UV6138 challenge. The resistance conferred by the gp96 derived from UV6138 cells was much greater (6 out of 7 mice) than the resistance conferred by the gp96 derived from the UV6139 against the UV6139SJ cells (2 out of 4 mice) (see U.S. Patent No. 5,837,251). These results indicate that administration of gp96 pre-



two UV-induced carcinomas immunized syngeneic mice from the type and that the resistance rendered was greater and more uniform immunogenic carcinoma cells.

Coadministration of one or more saponins along with complexes will elicit the desired immune response using reduced amounts of complexes as compared to gp96-peptide complexes administered alone (in the presence of saponin). Accordingly, the invention provides the advantage of the amount of an hsp or  $\alpha$ 2M-peptide complex required to elicit a response for prevention or treatment of cancer or infectious disease.

10

## 5.2. Treatment Modality Using gp96

Materials and Methods. The ability of gp96/antigen compositions to mediate therapy of pre-existing cancers is tested in mice. Mice are injected intradermally with  $10^7$  cells of the UV6139SJ carcinoma cells. Mice are kept under observation until the tumors become visible and palpable. Thereafter, mice in a first group receive no treatment. Mice in a second group receive every other day for a total of five injections, 6, 1, 0.6, or 0.1  $\mu$ g/mouse of a gp96-peptide complex derived from UV6139SJ carcinoma cells. Mice in a third group receive in a similar manner, a total of five injections of gp96-peptide complex derived from liver. Mice in a fourth group receive, in a similar manner, a total of five injections of a gp96-peptide complex derived from the UV6139SJ carcinoma cells, mixed with QS-21. The mice in the fifth group receive, in a similar manner, a total of five injections of a gp96-peptide complex derived from normal liver mixed with 20  $\mu$ g of QS-21. The mice in the sixth group receive 20 or 100  $\mu$ g QS-21 alone.

Results. In prior experiments, tumor growth, monitored by two perpendicular diameter measurements, was significantly retarded in mice treated with tumor-derived gp96 but not in mice treated with the liver-derived gp96 (see U.S. Patent No. 5,837,251). These results indicated a therapeutic effect of gp96-peptide complexes in the UV6139SJ carcinoma model. All mice in the sixth group showed tumor growth. A scrutiny of the kinetics of tumor growth in treated mice indicated that administration of tumor-derived gp96-peptide complex had an effect on tumor growth and that the effect appears to diminish after the gp96-peptide complex is terminated.

Coadministration of QS-21 with gp96-peptide complex derived from UV6139SJ carcinoma cells will result in a retardation of tumor growth with smaller amounts of gp96-peptide complex as compared to the group receiving

complex from UV6139SJ carcinoma cells without QS-21. The a alone, or gp96-peptide complex from normal liver plus QS-21, v significant difference as compared to the PBS control, or gp96-f normal liver, respectively.

5

### 5.3. Analogous Experiments Using $\alpha 2$

Use of  $\alpha 2$ M or other hsp in place of gp96 as des in both prevention of new tumors and retarded growth of establi amounts of the hsp or  $\alpha 2$ M when saponin adjuvant is present. A 10 expressed antigenic peptide - $\alpha 2$ M or -hsp fusion proteins can als immunosuppression of new tumors and retarded growth of estab reduced amounts of the hsp or  $\alpha 2$ M fusion proteins.

Accordingly, in analogous experiments, a saponin added to the aforementioned preparations and administered. The 15 compositions are used in experiments that parallel those describe compositions. Coadministration of QS-21 along with hsp- or  $\alpha 2$  a significant decrease in the amount of hsp- or  $\alpha 2$ M- complex re immunosuppression of new tumors and retarded growth of estab

20

The present invention is not to be limited in scope embodiments described herein. Indeed, various modifications of to those described herein will become apparent to those skilled in description. Such modifications are intended to fall within the s claims.

25

Various publications are cited herein, the disclosu incorporated by reference herein in their entireties.

30

35

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising a purified (hsp) and a saponin.  
5
2. The pharmaceutical composition of claim 1, further antigenic molecule.
3. The pharmaceutical composition of claim 2, wherei  
10 molecule is bound to the hsp to form an hsp-antigenic molecule co
4. The pharmaceutical composition of claim 3, wherei molecule is non-covalently bound to the hsp.
- 15 5. A pharmaceutical composition comprising a purified ( $\alpha$ 2M) and a saponin.
6. The pharmaceutical composition of claim 5, further antigenic molecule.  
20
7. The pharmaceutical composition of claim 6, wherei molecule is bound to the  $\alpha$ 2M to form an  $\alpha$ 2M-antigenic molecule
8. The pharmaceutical composition of claim 7, wherei  
25 molecule is non-covalently bound to the  $\alpha$ 2M.
9. The pharmaceutical composition of claim 2, wherei molecule is not bound to the saponin or to the hsp.
- 30 10. The pharmaceutical composition of claim 9, further antigenic molecule.
11. The pharmaceutical composition of claim 10, where molecule is bound to the hsp.  
35
12. The pharmaceutical composition of claim 11, where

molecule is covalently bound to the hsp to form an hsp-second anti

13. The pharmaceutical composition of claim 11, wherei  
molecule is non-covalently bound to the hsp.

5

14. The pharmaceutical composition of claim 10, further  
antigenic molecule.

15. The pharmaceutical composition of claim 14, wherei  
10 molecule is covalently bound to the saponin to form a saponin-third  
complex.

16. The pharmaceutical composition of claim 10, wherei  
molecule is covalently bound to the saponin.

15

17. The pharmaceutical composition of claim 2, wherei  
molecule is bound to the hsp to form an hsp-first antigenic molecule

18. The pharmaceutical composition of claim 17, where:  
20 molecule is covalently bound to the hsp.

19. The pharmaceutical composition of claim 17, where:  
molecule is non-covalently bound to the hsp.

20. The pharmaceutical composition of claim 17, further  
25 antigenic molecule covalently bound to the saponin to form a sapor  
molecule complex.

21. The pharmaceutical composition of claim 2, wherei  
30 molecule is covalently bound to the saponin to form a saponin-first  
complex.

22. The pharmaceutical composition of claim 2, wherei  
gp96, calreticulin, hsp110, grp170, PDI, or a mixture of two or moi

35

23. The pharmaceutical composition of claim 2, wherei

is at least 1 microgram.

24. The pharmaceutical composition of claim 23, where  
is 10 to 20 micrograms.

5

25. The pharmaceutical composition of claim 23, where  
is 20 to 100 micrograms.

26. The pharmaceutical composition of claim 23, where  
10 is 100 to 500 micrograms.

27. The pharmaceutical composition of claim 2, where  
least 0.1 microgram.

15 28. The pharmaceutical composition of claim 27, where  
least 1 microgram.

29. The pharmaceutical composition of claim 27, where  
no greater than 10 micrograms.

20

30. The pharmaceutical composition of claim 1 or 5, or  
QS-7, QS-21, QS-21-V1, or QS-21-V2.

31. The pharmaceutical composition of claim 30, where

25

32. The pharmaceutical composition of claim 1, where

33. The pharmaceutical composition of claim 2 or 6, where  
molecule displays antigenicity of a tumor-associated antigen.

30

34. The pharmaceutical composition of claim 33, where  
fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteo  
angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphoma  
synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhab  
35 carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate  
carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland ca

carcinoma, papillary carcinoma, papillary adenocarcinomas, cystic  
 medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma,  
 carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms  
 cancer, testicular tumor, lung carcinoma, small cell lung carcinoma,  
 5 epithelial carcinoma, glioma, astrocytoma, medulloblastoma, crani-  
 oependymoma, pinealoma, hemangioblastoma, acoustic neuroma, olfactory  
 meningioma, melanoma, neuroblastoma, retinoblastoma, leukemia,  
 lymphoma, multiple myeloma, Waldenström's macroglobulinemia.

10 35. The pharmaceutical composition of claim 2 or 6, where the  
 molecule displays antigenicity of an antigen of an agent of infectious

36. The pharmaceutical composition of claim 35, where the agent of infection  
 is a viral disease.

15

37. The pharmaceutical composition of claim 36, where the agent of infection  
 is type A, hepatitis type B, hepatitis type C, influenza, varicella, adenovirus,  
 type I (HSV-I), herpes simplex type II (HSV-II), rinderpest, rhinovirus,  
 rotavirus, respiratory syncytial virus, papilloma virus, papova virus,  
 20 echinovirus, arbovirus, hantavirus, coxsackie virus, mumps virus, measles  
 virus, polio virus, human immunodeficiency virus type I (HIV-I), and  
 human immunodeficiency virus type II (HIV-II).

38. The pharmaceutical composition of claim 35, where the agent of infection  
 25 is a bacterial disease.

39. The pharmaceutical composition of claim 38, where the agent of infection  
 is mycobacteria rickettsia, mycoplasma, neisseria and legionella.

30 40. The pharmaceutical composition of claim 35, where the agent of infection  
 is a protozoal disease.

41. The pharmaceutical composition of claim 40, where the agent of infection  
 is leishmania, kokzidioa, or trypanosoma.

35

42. The pharmaceutical composition of claim 2 or 6, where the agent of infection

molecule displays antigenicity of an antigen associated with a neuro

43. The pharmaceutical composition of claim 42, wherei  
molecule is  $\beta$ -amyloid or a fragment thereof, an oligomeric A $\beta$  cor  
5 thereof, an ApoE4-A $\beta$  complex or a fragment thereof, tau protein or  
mutant amyloid precursor protein or a fragment thereof, a mutant of  
fragment thereof,  $\alpha$ -synuclein or a fragment thereof, or a prion prot  
thereof.

10 44. The pharmaceutical composition of claim 42, wherei  
disorder is Alzheimer's Disease, age-related loss of cognitive functi  
Parkinson's disease, amyotrophic lateral sclerosis, Wilson's Diseas  
progressive supranuclear palsy, Guam disease, Lewy body dementi  
spongiform encephalopathy, Creutzfeldt-Jakob disease, a polygluta  
15 Huntington's disease, myotonic dystrophy, Freidrich's ataxia, ataxi  
syndrome, seizure disorders, epilepsy, chronic seizure disorder, strc  
cord trauma, AIDS dementia, alcoholism, autism, retinal ischemia,  
function disorder, hypertension, neuropsychiatric disorder, schizopl  
disorder.

20 45. The pharmaceutical composition of claim 11, where  
antigenic molecule complex is purified from a cancerous cell.

46. The pharmaceutical composition of claim 11, where  
25 antigenic molecule complex is purified from a cell infected with an  
disease.

47. The pharmaceutical composition of claim 17, where  
molecule complex is purified from a cancerous cell.

30 48. The pharmaceutical composition of claim 17, where  
molecule complex is purified from a cell infected with an agent of i

49. The pharmaceutical composition of claim 11, where  
35 antigenic molecule complex is prepared *in vitro*.

50. The pharmaceutical composition of claim 17, where molecule complex is prepared *in vitro*.

51. The pharmaceutical composition of claim 2, wherein  
5 fusion protein comprising the first antigenic molecule.

52. The pharmaceutical composition of claim 10, where fusion protein comprising the second antigenic molecule.

10 53. The pharmaceutical composition of claim 52, where molecule is not bound to the saponin or to the  $\alpha 2M$ .

54. The pharmaceutical composition of claim 53, further antigenic molecule.

15

55. The pharmaceutical composition of claim 54, where molecule is bound to the  $\alpha 2M$ .

56. The pharmaceutical composition of claim 55, where  
20 molecule is covalently bound to the  $\alpha 2M$  to form an  $\alpha 2M$ -second complex.

57. The pharmaceutical composition of claim 55, where molecule is non-covalently bound to the  $\alpha 2M$ .

25

58. The pharmaceutical composition of claim 54, further antigenic molecule.

59. The pharmaceutical composition of claim 58, where  
30 molecule is covalently bound to the saponin to form a saponin-thi complex.

60. The pharmaceutical composition of claim 54, where molecule is covalently bound to the saponin.

35

61. The pharmaceutical composition of claim 52, where



molecule is bound to the  $\alpha$ 2M to form an  $\alpha$ 2M-first antigenic mo-

62. The pharmaceutical composition of claim 61, when molecule is covalently bound to the  $\alpha$ 2M.

5

63. The pharmaceutical composition of claim 61, when molecule is non-covalently bound to the  $\alpha$ 2M.

64. The pharmaceutical composition of claim 61, further 10 antigenic molecule covalently bound to the saponin to form a saponin molecule complex.

65. The pharmaceutical composition of claim 52, when molecule is covalently bound to the saponin to form a saponin-fir 15 complex.

66. The pharmaceutical composition of claim 52, when is at least 1 microgram.

67. The pharmaceutical composition of claim 66, when 20 is 10 to 20 micrograms.

68. The pharmaceutical composition of claim 66, when is 20 to 100 micrograms.

25

69. The pharmaceutical composition of claim 66, when is 100 to 500 micrograms.

70. The pharmaceutical composition of claim 52, when 30 least 0.1 microgram.

71. The pharmaceutical composition of claim 70, when least 1 microgram.

72. The pharmaceutical composition of claim 70, when 35 no greater than 10 micrograms.

73. The pharmaceutical composition of claim 52, where

74. The pharmaceutical composition of claim 52, where  
5 recombinant.

75. The pharmaceutical composition of claim 52, where  
molecule displays antigenicity of a tumor-associated antigen.

10 76. The pharmaceutical composition of claim 52, where  
molecule displays antigenicity of an antigen of an agent of infection

77. The pharmaceutical composition of claim 52, where  
fusion protein comprising the first antigenic molecule.

15

78. The pharmaceutical composition of claim 54, where  
fusion protein comprising the second antigenic molecule.

79. A method of eliciting an immune response against  
20 infectious disease in an individual comprising administering to the  
a pharmaceutical composition comprising a purified hsp, a first antigen  
displays antigenicity of an antigen of said type of cancer or of an agent  
of infectious disease, and a saponin, which amount is effective to elicit  
the individual.

25

80. The method according to claim 79 in which the individual

81. The method according to claim 80 in which the major

30 82. The method according to claim 79, further comprising  
individual an effective amount of a biological response modifier selected  
consisting of interferon- $\alpha$ , interferon- $\gamma$ , interleukin-2, interleukin-6,  
tumor necrosis factor.

35 83. The method according to claim 79 in which the pharmaceutical  
composition is administered at weekly intervals.

84. The method according to claim 79 in which the pharmaceutical composition is administered intramuscularly, subcutaneously, intravenously, intradermally or mucosally.

5

85. A method of eliciting an immune response against an infectious disease in an individual comprising administering to the individual a pharmaceutical composition comprising a purified  $\alpha 2M$ , a first antigen displaying antigenicity of an antigen of said type of cancer or of an infectious disease, and a saponin, which amount is effective to elicit an immune response in the individual.

86. A method of treating or preventing cancer in an individual in whom such treatment or prevention is desired, comprising administering to the individual a pharmaceutical composition comprising a purified hsp, a first antigen displaying antigenicity of an antigen of said type of cancer, and a saponin, which amount is effective to treat or prevent cancer in the individual.

87. A method of treating or preventing cancer in an individual in whom such treatment or prevention is desired, comprising administering to the individual a pharmaceutical composition comprising a purified  $\alpha 2M$ , a first antigen displaying antigenicity of an antigen of said type of cancer, and a saponin, which amount is effective to treat or prevent cancer in the individual.

88. A method of treating or preventing an infectious disease in an individual in whom such treatment or prevention is desired, comprising administering to the individual an amount of a pharmaceutical composition comprising a purified antigenic molecule that displays antigenicity of an antigen of said infectious disease, and a saponin, which amount is effective to treat or prevent the infectious disease.

30

89. A method of treating or preventing a neurodegenerative disease in an individual in whom such treatment or prevention is desired, comprising administering to the individual an amount of a pharmaceutical composition comprising a first antigenic molecule that displays antigenicity of an antigen associated with a neurodegenerative or amyloid disease, and a saponin, which amount is effective to prevent cancer in the individual.

90. A method of treating or preventing a neurodegenerative disease in an individual in whom such treatment or prevention is desired, comprising administering to the individual an amount of a pharmaceutical composition comprising a first antigenic molecule that displays antigenicity of an antigen associated with a neurodegenerative or amyloid disease, and a saponin, which amount is effective to prevent cancer in the individual.

91. A method of treating or preventing an infectious disease in an individual in whom such treatment or prevention is desired, comprising administering to the individual an amount of a pharmaceutical composition comprising a purified antigenic molecule that displays antigenicity of an agent of said infectious disease, which amount is effective to treat or prevent the infectious disease.

92. A method of treating or preventing an autoimmune disease in an individual in whom such treatment or prevention is desired, comprising administering to the individual an amount of a pharmaceutical composition comprising a purified antigenic molecule that displays antigenicity of an agent of said autoimmune disease, which amount is effective to treat or prevent the autoimmune disease in the individual.

93. The method of claim 92, wherein the pharmaceutical composition comprises an antigenic molecule.

94. A method of treating or preventing an autoimmune disease in an individual in whom such treatment or prevention is desired, comprising administering to the individual an amount of a pharmaceutical composition comprising a purified antigenic molecule that displays antigenicity of an agent of said autoimmune disease, which amount is effective to treat or prevent the autoimmune disease in the individual.

95. The method of claim 94, wherein the pharmaceutical composition comprises an antigenic molecule.

96. A method of making a pharmaceutical composition comprising  $\alpha$ 2M, a first antigenic molecule that displays the antigenicity of an antigen associated with a cancer, infectious disease, neurodegenerative disease or autoimmune disease, and a saponin, said method comprising:

combining the saponin, the first antigenic molecule and  $\alpha$ 2M under conditions that produce a pharmaceutical composition.

97. The method of claim 96, wherein the saponin is covalently bound to the first antigenic molecule and then with the hsp or  $\alpha 2M$ .

98. The method of claim 97, wherein the saponin is covalently bound to the first antigenic molecule under conditions that do not promote covalent binding of the saponin and the first antigenic molecule.

99. The method of claim 97, wherein the saponin is covalently bound to the first antigenic molecule under conditions that promote covalent binding of the saponin and the first antigenic molecule.

100. The method of claim 97, wherein the hsp or  $\alpha 2M$  is covalently bound to the second antigenic molecule.

101. The method of claim 97, wherein the hsp or  $\alpha 2M$  is covalently bound to the second antigenic molecule.

102. The method of claim 101, wherein the hsp or  $\alpha 2M$  is covalently bound to the second antigenic molecule.

103. The method of claim 101, wherein the hsp or  $\alpha 2M$  is covalently bound to the second antigenic molecule.

104. The method of claim 97, wherein the hsp or  $\alpha 2M$  is covalently bound to the second antigenic molecule.

105. The method of claim 96, wherein the hsp or  $\alpha 2M$  is covalently bound to the second antigenic molecule and then with the saponin.

106. The method of claim 105, wherein the first antigenic molecule is covalently bound to the hsp or  $\alpha 2M$  under conditions that promote formation of a covalent complex of the first antigenic molecule and the hsp or  $\alpha 2M$ .

107. The method of claim 105, wherein the saponin is covalently bound to the second antigenic molecule.

## WO0211669

### Publication Title:

COMPOSITIONS COMPRISING HEAT SHOCK PROTEINS OR ALPHA(2)MACROGLOBULIN, ANTIGENIC MOLECULES AND SAPONINS, AND METHODS OF USE THEREOF

### Abstract:

The present invention relates to pharmaceutical compositions and methods for the prevention and treatment of autoimmune diseases, infectious diseases, neurodegenerative diseases, and primary and metastatic neoplastic diseases. In the practice of the invention, the compositions are employed comprising: (a) a heat shock protein (hsp) or an alpha(2)macroglobulin (alpha 2M); (b) a saponin; and, optionally, (c) an antigenic molecule. The antigenic molecule displays the antigenicity of an antigen of: (a) a cell that elicits an autoimmune response; (b) an agent of an infectious disease; (c) a cancerous cell; or (d) a cell or structure associated with a neurodegenerative or amyloid disease. The hsps that can be used in the practice of the invention include but are not limited to hsp70, hsp90, gp96, calreticulin, hsp 110, gp170, and PDI, alone or in combination with each other. The antigenic molecule can be covalently or noncovalently bound to the hsp or alpha 2M, free in solution, and/or covalently bound to the saponin. The compositions of the invention can be administered alone or in combination with the administration of antigen presenting cells sensitized with an hsp- or alpha 2M-antigenic molecule complex.

-----  
Data supplied from the esp@cenet database - <http://ep.espacenet.com>

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ ~~FADED~~ TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**